

Searches for long-lived particles at LHCb



Marcin Chrząszcz
mchrzasz@cern.ch




University of
Zurich^{UZH}

SUSY 2015, Tahoe City, 23-29 August, 2015



Why long living particles?

- We all know here that the SM is incomplete.
- Unfortunately we do not know what is the scale of NP.
- NP still can come from the Higgs sector \Rightarrow not all properties are yet constrained.





- There is a long list of theoretical models that predict the existence of new particles that couple to the SM sector by mixing with the Higgs.
- Inflaton, axion-like, dark matter mediator models also predict the new boson to be light.
- SUSY models also can have create stable long living particles like \tilde{q} , $\tilde{\ell}$.

LHCb detector - tracking




- Excellent Impact Parameter (IP) resolution ($20 \mu\text{m}$).
⇒ Identify secondary vertices from heavy flavour decays
- Proper time resolution $\sim 40 \text{ fs}$.
⇒ Good separation of primary and secondary vertices.
- Excellent momentum ($\delta p/p \sim 0.4 - 0.6\%$) and inv. mass resolution.
⇒ Low combinatorial background.

LHCb detector - particle identification



- Excellent Muon identification $\epsilon_{\mu \rightarrow \mu} \sim 97\%$, $\epsilon_{\pi \rightarrow \mu} \sim 1 - 3\%$
- Good $K - \pi$ separation via RICH detectors, $\epsilon_{K \rightarrow K} \sim 95\%$,
 $\epsilon_{\pi \rightarrow K} \sim 5\%$.
⇒ Reject peaking backgrounds.
- High trigger efficiencies, low momentum thresholds. Muons:
 $p_T > 1.76 \text{ GeV}$ at L0, $p_T > 1.0 \text{ GeV}$ at HLT1,
 $B \rightarrow J/\psi X$: Trigger $\sim 90\%$.

Data taken by LHCb




- In 2011 and 2012 LHCb has gather 3 fb^{-1} of pp collisions.

$B \rightarrow K^* \chi(\mu\mu)$ search

- Search for displaced di-muon vertex coming from B meson.

$$B^0 \rightarrow K^* \chi(\mu^- \mu^+)$$



- If χ mixes with the Higgs and it is light:
 - $\Gamma(K \rightarrow \pi \chi) \propto m_t^4 \lambda^5$
 - $\Gamma(D \rightarrow \pi \chi) \propto m_b^4 \lambda^5$
 - $\Gamma(D \rightarrow K \chi) \propto m_t^4 \lambda^2$
- In addition $K^* \rightarrow K \pi$ helps in vertex reconstruction.
- High $\mathcal{B}(\chi \rightarrow \mu^- \mu^+)$.

$B \rightarrow K^* \chi(\mu\mu)$ motivation

Possible models:

1. Inflaton: [Phys.Lett. B736 \(2014\) 494](#)

- $\tau_\chi = 10^{-8} - 10^{-10} \text{ s}$
- $m_\chi \mathcal{O}(1 \text{ GeV})$
- $\mathcal{B}(B \rightarrow K \chi) \sim 10^{-6}$
- effective couplings to SM particles:
 - $g_Y \frac{m_f}{v_{EW}}, g_Y = \sin \theta$

2. Axion portal: [Phys.Rev.D81:034001,2010](#)

- Prompt decay.
- Large allowed masses.
- Axion decay constant: $f_\chi \sim 1 - 3 \text{ TeV}$
 - Coupling $\propto \frac{m_f}{f_\chi}$.


All those particles have width much smaller than resolution of LHCb detector.

Signal properties

⇒ Depending on the coupling of the hidden sector we can identify two lifetime regimes:


Long lifetime (> 0.2 ps)

- Inflaton [JHEP 1005:010](#)
- Displaced vertex.
- Almost background free.
- Lower reconstruction efficiency.



Long lifetime (≤ 0.2 ps)

- Dark matter mediator [Phys. Lett. B727](#)
- Axion [Phys.Rev.D81](#)
- Prompt decay.
- Contaminated via Sm decay.



Selection

- Trigger on muons.
- Multivariate selection: μ BDT <http://arxiv.org/abs/1305.7248>
 - μ BDT ensures flat efficiency in lifetime of χ .
- Optimized on Punzi figure-of-merit:




$$P_a = \frac{S}{\frac{5}{2} + \sqrt{B}},$$

with S and B are signal and background yields.


- Factorize lifetime into two components: $\mathcal{L} = \mathcal{L}^{\text{prompt}} \otimes \mathcal{L}^{\text{displaced}}$
 - Prompt: $\tau < 3\sigma_\tau$
↳ SM background of $B^0 \rightarrow K^* \mu^- \mu^+$
 - Displaced: $\tau > 3\sigma_\tau$
↳ Almost background free.

Search strategy

- B^0 mass constrained.
- Di-muon mass resolution $\sigma_m = 1 - 7$ MeV.
- Scan m_{test} in steps of $0.5 \sigma_m$.
 - **Wide resonances** can't affect the search.
 - **Narrows resonances** one we veto.
- Calculations performed in each m_{test} window.
- A global p-value is assigned from minimum local p-value observed.




Results



- ⇒ Grey regions correspond to vetoed regions where narrow resonances are expected.
- ⇒ Largest deviation seen in $m_\chi = 253$ MeV.
- Not statistically significant: local p-value = 0.2.
- ⇒ LHCb-PAPER-2015-036 in preparation.

Branching fraction exclusion limit



⇒ No deviations from background only hypothesis is observed.


- We set a 95% CL upper limit as function of mass and lifetime of the new particle (in the LHCb accessible range).
- Lower lifetimes have better limit due to higher reconstruction efficiency.

Benchmark models

⇒ Interpretation of the results in to specific models:


(Specific) inflaton model

[LHCb-PAPER-2015-036 in preparation]



Include 3 sterile neutrinos N_i

Axion portal



MSSM-like two Higgs doublet model.

Long living charged particles

⇒ Long living particles can also be produced in the PV.



- This kind of particles would be produced in relatively low velocities and could be identified by their time-of-flight, dE/dx or in Cherenkov detectors.

⇒ LHCb performed a search for long living $\tilde{\tau}$ particles.

⇒ $\tilde{\tau}^+\tilde{\tau}^-$ produced by Drell-Yan process.

$\tilde{\tau}$ analysis strategy


- ⇒ Search performed $\tilde{\tau}$ in mass range of 124 – 309 GeV.
- ⇒ After the loose preselection to reduce normal Drell-Yan production.



- ⇒ After the preselection a Neural Network is trained based on Cherenkov detectors to calculate to further suppress the remaining background.

$\tilde{\tau}$ results

- No significant signal yield has been observed.
- Upper limit has been set.



Conclusion

- A search for a dark boson in the decay channel $B^0 \rightarrow K^* \mu^- \mu^+$ has been presented.
 - No deviations from SM observed.
- Results are the most constraining exclusion limit on the process.
- LHCb is suited for search for long living particles.
- Stay tuned, more searches like this are on they way.

