

# $\Lambda_c^+ \rightarrow p\mu\mu$

## Status Update and Plans for future



Marcin Chrzaszcz



University of  
Zurich<sup>UZH</sup>

on behalf of the  $\Lambda_c^+ \rightarrow p\mu\mu$  team:

T.Lesiak(IFJ, Krakow), B.Nowak(IFJ, Krakow), M.Witek (IFJ, Krakow),  
L.Pescatore(EPFL, Lausanne), M.Stamenkovic(EPFL, Lausanne),  
M.Martinelli(EPFL, Lausanne)

Analysis and Software Week, CERN  
April, 2017

# Topics covered in this presentation


1. Physics of  $\Lambda_c^+ \rightarrow p\mu\mu$
2. Pre-Selection.
3. MVA selection.
4. PID.
5. Normalization.
6. Systematics.
7. Expected limits.
8. Run2 extensions.

## Yellow pages

- ⇒ Review started on 31.03.2017.
- ⇒ Reviewers: Tom Blake, Harry Cliff; many thanks for refereeing!
- ⇒ Twiki:  
<https://twiki.cern.ch/twiki/bin/view/LHCbPhysics/Lc2PMuMu>
- ⇒ The newest version of the ANA note:  
CLIC

# Physics of $\Lambda_c^+ \rightarrow p\mu\mu$

$\Rightarrow \Lambda_c^+ \rightarrow p\mu\mu$  is a FCNC in the charm sector:




$\Rightarrow$  Current experimental situation:

$\Rightarrow$  SM prediction:

- Short distance  
 $Br \sim \mathcal{O}(10^{-8})$
- Long distance  
 $Br \sim \mathcal{O}(10^{-6})$
- Expected to improve by  $\mathcal{O}(10^2)$

- $Br(\Lambda_c^+ \rightarrow p\mu\mu) < 4.4 \times 10^{-5}$  at 90 %CL  
arXiv:1107.4465 (BaBar)



# Strategy

- ⇒ We follow the strategy of previous analysis:  $\tau \rightarrow \mu\mu\mu$  and  $\tau \rightarrow p\mu\mu$ .
- ⇒ Analysis based on 2011 and 2012 data sets.
- ⇒ Blind the signal window:  $|m_{p\mu\mu} - m_{\Lambda_c^+}^{PDG}| < 40$  MeV
- ⇒ We start from stripping and loose pre-selection.
- ⇒ MVA:
  - Signal MC.
  - Background side-bands.
- ⇒ k-Folding technique applied.
- ⇒ Two BDT are used:
  - BDT1 to first clean up the sample.
  - BDT2 to further increase the sensitivity.
- ⇒ Final 3D optimization: (BDT2, ProbNNp, ProbNNmu).
- ⇒ Calculate the UL with  $CL_s$ .

# Trigger

⇒ We decided to base the analysis on muon triggers:

- L0
  - Lambda\_cplus\_L0MuonDecision\_TOS
  - Lambda\_cplus\_L0DiMuonDecision\_TOS
- HLT1
  - Lambda\_cplus\_Hlt1TrackMuonDecision\_TOS
  - Lambda\_cplus\_Hlt1DiMuonLowMassDecision\_TOS
  - Lambda\_cplus\_Hlt1TrackAllL0Decision\_TOS
- HLT2
  - Lambda\_cplus\_Hlt2CharmHadD2HHHDecision\_TOS;
  - Lambda\_cplus\_Hlt2DiMuonDetachedDecision\_TOS;
  - Lambda\_cplus\_Hlt2CharmSemilep3bodyD2KMuMuDecision\_TOS;
  - Lambda\_cplus\_Hlt2CharmSemilepD2HMuMuDecision\_TOS;

# Stripping

StrippingTau23MuTau2PMuMuLine	
Condition	$\Lambda_c^+ \rightarrow p \mu \mu$
$\mu^\pm$ and $p_T$	$> 300 \text{ MeV}/c$
Track $\chi^2/\text{ndf}$	$< 3$
IP $\chi^2/\text{ndf}$	$> 9$
PID $\mu^\pm$	PIDmu > -5 and (PIDmu - PIDK) > 0
PID p	PIDp > 10
$\Lambda_c^+$	
$\Delta m$	$< 150 \text{ MeV}/c^2$
Vertex $\chi^2$	$< 15$
IP $\chi^2$	$< 225$
$c\tau$	$> 100 \mu\text{m}$
Lifetime fit $\chi^2$	$< 225$

⇒ In Run2 we have a dedicated stripping/HLT2 lines for  $\mu, e$  lepton flavours.


## Futher preselection

Common cuts
$m_{\mu\mu} > 250 \text{ MeV}/c^2$
proton $ProbNNp > 0.1$
$\mu^+, \mu^-$ $ProbNNmu > 0.1$
$10 \text{ GeV}/c < p_{\text{proton}} < 100 \text{ GeV}/c$
Signal channel
$ m_{\mu\mu} - m_\omega  > 40 \text{ MeV}/c^2$
$ m_{\mu\mu} - m_\phi  > 40 \text{ MeV}/c^2$
Normalization channel
$ m_{\mu\mu} - m_\phi  < 35 \text{ MeV}/c^2$



## MVA Selection 1/2

⇒ The BDT1 uses a small set of available variables related to  $\Lambda_c^+$  candidate:


- Lambda\_cplus\_IP\_OWNPV
- Lambda\_cplus\_IPCHI2\_OWNPV
- TMath :: Exp( $-1000 * \text{Lambda\_cplus\_TAU}$ )
- Lambda\_cplus\_ENDVERTEX\_CHI2
- Lambda\_cplus\_PT
- Lambda\_cplus\_FD\_OWNPV
- Lambda\_cplus\_FDCHI2\_OWNPV



## MVA Selection 2/2



⇒ We choose a loose cut ( $\text{BDT1} > -0.1$ ) to clean up the sample:



$$\Lambda_c \rightarrow p\phi(\mu\mu)$$

# Normalization

⇒  $\Lambda_c \rightarrow p\phi(\mu\mu)$ :

- Same final state!
- Most of the systematics cancel in the ratio.
- Kinematics difference will only remain.
- Low Br:  $Br(\Lambda_c \rightarrow p\phi(\mu\mu)) = (2.98 \pm 0.63) \times 10^{-7}$




⇒  $\Lambda_c \rightarrow p\pi\pi$ :

- Different final state!
- The systematics will not cancel in the ratio.
- Need to understand the  $\pi\pi$  spectrum.
- High branching fraction:  
 $Br(\Lambda_c \rightarrow p\pi\pi) = (4.3 \pm 2.3) \times 10^{-3}$

We have chosen the  $\Lambda_c \rightarrow p\phi(\mu\mu)$  as normalization channel.

## MVA Selection II

- Added variables related to the daughter tracks.






⇒ The BDT was checked against the correlation with mass on MC background.

⇒ All cross-checks passed.


⇒ The PID in this analysis is done using re sampling the PID distributions.

- PIDCalib for muons does not cover the low  $p_T$  muons (10 %) of the sample.
- We used the  $D_s \rightarrow \pi\phi(\mu\mu)$ .
- The same procedure was used in the different analysis with this problem.
- The sample is currently being included to the standard sample from the PID WG.






# Optimization

- ⇒ Optimization was performed on a TOY MC sample.
- ⇒ The toys were generated using PDF from signal MC and sideband sample.
- ⇒ Optimization was done on grid of points, using 100 TOYs per point.
- ⇒  $CL_s$  was used as FOM.



Variable	Cut
BDT2	> 0.0
ProbNNp	> 0.68
ProbNNmu	> 0.38



## Peaking backgrounds 1/2

⇒ There are several sources of peaking background:

Resonance	$\text{BF}(\Lambda_c^+ \rightarrow p X)$	$\text{BF}(X \rightarrow \mu\mu)$	Total BF
$\eta$	-	$(5.8 \pm 0.8) \times 10^{-6}$	-
$\rho$	-	$(4.55 \pm 0.28) \times 10^{-5}$	-
$\omega$	-	$(9.0 \pm 3.1) \times 10^{-5}$	-
$\phi$	$(1.04 \pm 0.21) \times 10^{-3}$	$(2.87 \pm 0.19) \times 10^{-4}$	$(2.98 \pm 0.63) \times 10^{-7}$



Resonance	$\text{BF}(\Lambda_c^+ \rightarrow p X)$	$\text{BF}(X \rightarrow \mu\mu\gamma)$	Total BF
$\eta$	-	$(3.1 \pm 0.4) \times 10^{-4}$	-
$\eta'$	-	$(1.08 \pm 0.27) \times 10^{-4}$	-

⇒ Unfortunately not all of the BF are known...

⇒ We took the adequate decay of D mesons. We ended up with  $\text{BF} \mathcal{O}(10^{-9})$  for not vetoed decays, which is much below our sensitivity (see further slides).

## Peaking backgrounds 2/2

- ⇒ The other peaking background is a harmonic decay  $\Lambda_c^+ \rightarrow p\pi\pi$ .
- ⇒ Estimated from MC sample
- ⇒ Used the resampled PID response.
- ⇒ Observed number of events in the signal window.



- ⇒ Estimated:  $N_{\Lambda_c^+ \rightarrow p\pi\pi} = 1.96 \pm 1.13$
- ⇒ Took into account in background estimation.

# Normalization

⇒ Master equation:


$$\frac{Br(\Lambda_c \rightarrow p\mu\mu)}{Br(\Lambda_c \rightarrow p\phi(\mu\mu))} = \frac{\epsilon_{\text{norm}}^{\text{TOT}}}{\epsilon_{\text{sig}}^{\text{TOT}}} \times \frac{N_{\text{sig}}}{N_{\text{norm}}},$$

where

$$\frac{\epsilon_{\text{norm}}^{\text{TOT}}}{\epsilon_{\text{sig}}^{\text{TOT}}} = \frac{\epsilon_{\text{norm}}^{\text{STRIP}}}{\epsilon_{\text{sig}}^{\text{STRIP}}} \times \frac{\epsilon_{\text{norm}}^{\text{COMM}}}{\epsilon_{\text{sig}}^{\text{COMM}}} \times \frac{\epsilon_{\text{norm}}^{\text{SPEC}}}{\epsilon_{\text{sig}}^{\text{SPEC}}}$$

⇒ Signal window divided in 6 equal bins ( $7 \text{ MeV}/c^2$ )



⇒ Many of the ratios close to one:



Uncertainty source	Value
Efficiency ratio $R_{strip}$ (statistical)	0.2 %
Efficiency ratio $R_{comm}$ (statistical)	3.37 %
Efficiency ratio $R_{comm}$ (BDT2 cut)	0.4 %
Efficiency ratio $R_{comm}$ (PIDCalib samples)	0.71 %
Width of the signal peak	0.55 %
Yield of normalization channel	11.8 %
Dedicated PID resampling	0.26 %
$\Lambda_c \rightarrow p\phi(\mu\mu)$	21.5 %
Variation of signal decay model	15.3 %

# Expected limits

⇒ Putting all together one gets:



The expected limits:

$$Br(\Lambda_c \rightarrow p\mu\mu) < 5.9 \times 10^{-8} \text{ at 90% CL}$$

⇒ The RC started looking at the ANA note.


# Run 2 plans

⇒ We already started working on Run2 analysis.


⇒ The program is expanding:

- $Br(\Lambda_c^+ \rightarrow p\phi)$
- $Br(\Lambda_c^+ \rightarrow p\mu\mu)$
- $R(\Lambda_c^+) = \frac{Br(\Lambda_c^+ \rightarrow p\mu\mu)}{Br(\Lambda_c^+ \rightarrow p\mu e)}$
- LFV:  $\Lambda_c \rightarrow p\mu e$
- and maybe more ideas?

⇒ Prompt:



⇒ Semileptonics:



⇒  $\Lambda_c^+$  is a exciting system that is not fully explored!

⇒ We have a rich physics program to be studied with Run2 data.

# Backup

---