

Submitting results to HepData



Marcin Chrzaszcz
mchrzasz@cern.ch



**University of
Zurich**^{UZH}

With M.Bettler, A.Puig, A. Grecu

Zurich meeting, CERN
November 24, 2016

About Submission Help Sign in

HEPData
High Energy Physics Data Repository

This new site is still under development. Please continue Submitting Data using the old site at <http://hepdata.cedar.ac.uk>.

Search on 8336 publications and 66717 data Tables.

Search for papers, authors, experiments, reactions Search Advanced

e.g. reaction $PP \rightarrow LQ, LQ, X$, site from "sheldon collisions", collaboration in LHC or DG.

Data from the LHC

ATLAS [View Data](#)

ALICE [View Data](#)

CMS [View Data](#)

LHCb [View Data](#)

Recently Updated Submissions - View all

ALICE Jet-like correlations with neutral pion triggers in pp and central Pb-Pb collisions at 2.76 TeV
Adam, Jaroslav et al

CMS Distributions of Topological Observables in Inclusive Three- and Four-Jet Events in pp Collisions at $\sqrt{s}=7$ TeV
Khachatryan, Vardan et al

LHCb Measurement of the τ_b [1,5] production cross-section in proton-proton collisions via the decay $\tau_b(1,5) \rightarrow \mu \mu$
Aaij, Rolf et al

This website uses cookies to ensure you get the best experience on our website [More info](#) [Get IT!](#)

Why do we need HepData?

Download All

Filter 83 data tables

Table 1

Data from Appendix A, Table 3
10.17182/hepdata.74247.v1/1
CP-averaged angular observables evaluated by the unbinned maximum likelihood fit.

Table 2

Data from Appendix A, Table 4
10.17182/hepdata.74247.v1/2
CP-averaged angular observables evaluated by the unbinned maximum likelihood fit. The first uncertainties are statistical and the second systematic.

Table 3

Data from Appendix A, Table 5
10.17182/hepdata.74247.v1/3
CP-symmetric angular observables evaluated by the unbinned maximum likelihood fit. The first uncertainties are statistical and the second systematic.

Table 4

Data from Appendix A, Table 6
10.17182/hepdata.74247.v1/4
Optimized angular observables evaluated by the unbinned maximum likelihood fit. The first uncertainties are statistical and the second systematic.

Table 5

Data from Appendix A, Table 7
10.17182/hepdata.74247.v1/5
CP-asymmetric observables evaluated using the method of moments. The first uncertainties are statistical and the second systematic.

Table 6

Data from Appendix A, Table 8
10.17182/hepdata.74247.v1/6
CP-symmetries evaluated using the method of moments. The first uncertainties are statistical and the second systematic.

Table 7

Data from Appendix A, Table 9
10.17182/hepdata.74247.v1/7
Optimized observables evaluated using the method of moments. The first uncertainties are statistical and the second systematic.

Table 2 10.17182/hepdata.74247.v1/2

CP-averaged angular observables evaluated by the unbinned maximum likelihood fit. The first uncertainties are statistical and the second systematic.

cmenergies		observables							
7000.0-8000.0		FCL							
RE	PP →→ BD < K*(B92) < K+ Fl- → MU+ MU- > X								
SQRT(S)	7000.0 GeV								
SQRT(S)	8000.0 GeV								
q ² [GeV ²]	Fl	S ₁	S ₂	S ₃	A _{FB}	S ₇	S ₈	S ₉	
0.10-0.98	0.263 stat sys	-0.036 stat sys	0.082 stat sys	0.17 stat sys	-0.003 stat sys	0.015 stat sys	0.079 stat sys	-0.083 stat sys	
1.10-2.50	0.66 stat sys	-0.077 stat sys	-0.077 stat sys	0.137 stat sys	-0.191 stat sys	-0.219 stat sys	-0.098 stat sys	-0.119 stat sys	
2.50-4.00	0.876 stat sys	0.035 stat sys	-0.234 stat sys	-0.022 stat sys	-0.138 stat sys	0.068 stat sys	0.03 stat sys	-0.092 stat sys	
4.00-6.00	0.611 stat sys	0.035 stat sys	-0.219 stat sys	-0.146 stat sys	0.025 stat sys	-0.016 stat sys	0.167 stat sys	-0.032 stat sys	
6.00-8.00	0.579 stat sys	-0.042 stat sys	-0.296 stat sys	-0.249 stat sys	0.152 stat sys	-0.047 stat sys	-0.085 stat sys	-0.024 stat sys	
11.00-12.50	0.493 stat sys	-0.189 stat sys	-0.283 stat sys	-0.327 stat sys	0.318 stat sys	-0.141 stat sys	-0.007 stat sys	-0.004 stat sys	
15.00-17.00	0.349 stat sys	-0.142 stat sys	-0.321 stat sys	-0.316 stat sys	0.411 stat sys	0.061 stat sys	0.003 stat sys	-0.019 stat sys	
17.00-19.00	0.354 stat sys	-0.188 stat sys	-0.266 stat sys	-0.323 stat sys	0.305 stat sys	0.044 stat sys	0.013 stat sys	-0.094 stat sys	


http://www.hepdata.net

YAML
YODA
ROOT
CSV

reactions
PP →→ BD + X

- phrases
- Inclusive
 - Polarization
 - Proton-Proton Scattering
 - Strange Production
 - Muon Production

Visualize



Sum errors Log Scale (X)

Deselect variables or hide different error bars by clicking on them.

Variables

- Fl Sunned error
- S₁ Sunned error
- S₂ Sunned error
- S₃ Sunned error
- S₇ Sunned error
- A_{FB} Sunned error

The challenge

- ⇒ The $B \rightarrow K^* \mu \mu$ analysis has produced over 80 tables with results and correlation tables...
- ⇒ Theorists need all the correlation tables to make the global fit.
- ⇒ Reading the numbers from our tex files from all the theories groups is not really a nice way to do it:



- ⇒ Many thanks for Alex Grecu for help understanding how the HepData works!

Automatic scripts

⇒ Feel free to use, share and improve the scripts:

<https://gitlab.cern.ch/LHCb-RD/tools-hepdata>

<https://git.physik.uzh.ch/gitbucket/mchrzasz/HepData>

<http://www.physik.uzh.ch/~mchrzasz/HepData/>

Example:

- ⇒ The scripts I show here are not 100 % plug and play.
- ⇒ Each table in latex is a bit different and needs some special modification.
- ⇒ But to modify the scripts and apply them to your case should not be much work.

Example:

Table 4: CP -averaged angular observables evaluated by the unbinned maximum likelihood fit. The first uncertainties are statistical and the second systematic.

	$0.10 < q^2 < 0.98 \text{ GeV}^2/c^4$	$1.1 < q^2 < 2.5 \text{ GeV}^2/c^4$	$2.5 < q^2 < 4.0 \text{ GeV}^2/c^4$
F_L	$0.263^{+0.045}_{-0.044} \pm 0.017$	$0.660^{+0.083}_{-0.077} \pm 0.022$	$0.876^{+0.109}_{-0.097} \pm 0.017$
S_3	$-0.036^{+0.063}_{-0.063} \pm 0.005$	$-0.077^{+0.087}_{-0.105} \pm 0.005$	$0.035^{+0.098}_{-0.089} \pm 0.007$
S_4	$0.082^{+0.068}_{-0.069} \pm 0.009$	$-0.077^{+0.111}_{-0.113} \pm 0.005$	$-0.234^{+0.127}_{-0.144} \pm 0.006$
S_5	$0.170^{+0.059}_{-0.058} \pm 0.018$	$0.137^{+0.099}_{-0.094} \pm 0.009$	$-0.022^{+0.110}_{-0.103} \pm 0.008$
A_{FB}	$-0.003^{+0.055}_{-0.057} \pm 0.009$	$-0.191^{+0.068}_{-0.080} \pm 0.012$	$-0.118^{+0.082}_{-0.090} \pm 0.007$
S_7	$0.015^{+0.059}_{-0.059} \pm 0.006$	$-0.219^{+0.094}_{-0.104} \pm 0.004$	$0.068^{+0.120}_{-0.112} \pm 0.005$
S_8	$0.079^{+0.076}_{-0.075} \pm 0.007$	$-0.098^{+0.108}_{-0.123} \pm 0.005$	$0.030^{+0.129}_{-0.131} \pm 0.006$
S_9	$-0.083^{+0.058}_{-0.057} \pm 0.004$	$-0.119^{+0.087}_{-0.104} \pm 0.005$	$-0.092^{+0.105}_{-0.125} \pm 0.007$

	$4.0 < q^2 < 6.0 \text{ GeV}^2/c^4$	$6.0 < q^2 < 8.0 \text{ GeV}^2/c^4$	$11.0 < q^2 < 12.5 \text{ GeV}^2/c^4$
F_L	$0.611^{+0.052}_{-0.053} \pm 0.017$	$0.579^{+0.046}_{-0.046} \pm 0.015$	$0.493^{+0.049}_{-0.047} \pm 0.013$
S_3	$0.035^{+0.069}_{-0.068} \pm 0.007$	$-0.042^{+0.058}_{-0.059} \pm 0.011$	$-0.189^{+0.054}_{-0.058} \pm 0.005$
S_4	$-0.219^{+0.086}_{-0.084} \pm 0.008$	$-0.296^{+0.063}_{-0.067} \pm 0.011$	$-0.283^{+0.084}_{-0.095} \pm 0.009$
S_5	$-0.146^{+0.077}_{-0.075} \pm 0.011$	$-0.249^{+0.059}_{-0.060} \pm 0.012$	$-0.327^{+0.076}_{-0.079} \pm 0.009$
A_{FB}	$0.025^{+0.051}_{-0.052} \pm 0.004$	$0.152^{+0.041}_{-0.040} \pm 0.008$	$0.318^{+0.044}_{-0.040} \pm 0.009$
S_7	$-0.016^{+0.081}_{-0.080} \pm 0.004$	$-0.047^{+0.068}_{-0.066} \pm 0.003$	$-0.141^{+0.072}_{-0.074} \pm 0.005$
S_8	$0.167^{+0.094}_{-0.091} \pm 0.004$	$-0.085^{+0.072}_{-0.070} \pm 0.006$	$-0.007^{+0.070}_{-0.072} \pm 0.005$
S_9	$-0.032^{+0.071}_{-0.071} \pm 0.004$	$-0.024^{+0.059}_{-0.060} \pm 0.005$	$-0.004^{+0.070}_{-0.073} \pm 0.006$

	$15.0 < q^2 < 17.0 \text{ GeV}^2/c^4$	$17.0 < q^2 < 19.0 \text{ GeV}^2/c^4$
F_L	$0.349^{+0.039}_{-0.039} \pm 0.009$	$0.354^{+0.049}_{-0.048} \pm 0.025$
S_3	$-0.142^{+0.044}_{-0.049} \pm 0.007$	$-0.188^{+0.074}_{-0.084} \pm 0.017$
S_4	$-0.321^{+0.055}_{-0.074} \pm 0.007$	$-0.266^{+0.063}_{-0.072} \pm 0.010$
S_5	$-0.316^{+0.051}_{-0.057} \pm 0.009$	$-0.323^{+0.063}_{-0.072} \pm 0.009$
A_{FB}	$0.411^{+0.041}_{-0.037} \pm 0.008$	$0.305^{+0.049}_{-0.048} \pm 0.013$
S_7	$0.061^{+0.058}_{-0.058} \pm 0.005$	$0.044^{+0.073}_{-0.072} \pm 0.013$
S_8	$0.003^{+0.061}_{-0.061} \pm 0.003$	$0.013^{+0.071}_{-0.070} \pm 0.005$
S_9	$-0.019^{+0.054}_{-0.056} \pm 0.004$	$-0.094^{+0.065}_{-0.067} \pm 0.004$

Example:

RE P P → B0 < K*(892) < K+ Pi- → MU+ MU- → X									
SQRT(S) 7000.0 GeV									
SQRT(S) 8000.0 GeV									
χ^2 IN GeV^2	F_L	S_2	S_4	S_5	A_{FB}	S_7	S_8	S_9	
0.10 - 0.98	0.263 ± 0.045 - 0.044 (stat) ± 0.017 (sys)	-0.036 ± 0.063 (stat) ± 0.005 (sys)	0.082 ± 0.066 - 0.069 (stat) ± 0.009 (sys)	0.170 ± 0.059 - 0.058 (stat) ± 0.018 (sys)	-0.003 ± 0.058 - 0.057 (stat) ± 0.009 (sys)	0.015 ± 0.069 (stat) ± 0.006 (sys)	0.079 ± 0.076 - 0.075 (stat) ± 0.007 (sys)	-0.083 ± 0.058 - 0.057 (stat) ± 0.004 (sys)	
1.1 - 2.5	0.660 ± 0.083 - 0.077 (stat) ± 0.022 (sys)	-0.077 ± 0.087 - 0.105 (stat) ± 0.005 (sys)	-0.077 ± 0.111 - 0.113 (stat) ± 0.005 (sys)	0.137 ± 0.099 - 0.094 (stat) ± 0.009 (sys)	-0.191 ± 0.068 - 0.080 (stat) ± 0.012 (sys)	-0.219 ± 0.094 - 0.104 (stat) ± 0.004 (sys)	-0.098 ± 0.108 - 0.123 (stat) ± 0.005 (sys)	-0.119 ± 0.087 - 0.104 (stat) ± 0.005 (sys)	
2.5 - 4.0	0.876 ± 0.109 - 0.097 (stat) ± 0.017 (sys)	0.035 ± 0.098 - 0.089 (stat) ± 0.007 (sys)	-0.234 ± 0.127 - 0.144 (stat) ± 0.006 (sys)	-0.022 ± 0.110 - 0.103 (stat) ± 0.000 (sys)	-0.118 ± 0.082 - 0.090 (stat) ± 0.007 (sys)	0.068 ± 0.120 - 0.112 (stat) ± 0.005 (sys)	0.030 ± 0.129 - 0.131 (stat) ± 0.006 (sys)	-0.092 ± 0.105 - 0.125 (stat) ± 0.007 (sys)	
4.0 - 6.0	0.611 ± 0.052 - 0.053 (stat) ± 0.017 (sys)	0.035 ± 0.068 - 0.068 (stat) ± 0.007 (sys)	-0.219 ± 0.086 - 0.084 (stat) ± 0.008 (sys)	-0.146 ± 0.077 - 0.078 (stat) ± 0.011 (sys)	0.025 ± 0.051 - 0.052 (stat) ± 0.004 (sys)	-0.016 ± 0.081 - 0.080 (stat) ± 0.004 (sys)	0.167 ± 0.094 - 0.091 (stat) ± 0.004 (sys)	-0.032 ± 0.071 (stat) ± 0.004 (sys)	
6.0 - 8.0	0.579 ± 0.046 (stat) ± 0.015 (sys)	-0.042 ± 0.058 - 0.059 (stat) ± 0.011 (sys)	-0.296 ± 0.063 - 0.067 (stat) ± 0.011 (sys)	-0.249 ± 0.059 - 0.060 (stat) ± 0.012 (sys)	0.152 ± 0.041 - 0.040 (stat) ± 0.008 (sys)	-0.047 ± 0.068 - 0.066 (stat) ± 0.003 (sys)	-0.085 ± 0.072 - 0.070 (stat) ± 0.006 (sys)	-0.024 ± 0.059 - 0.060 (stat) ± 0.005 (sys)	
11.0 - 12.5	0.493 ± 0.049 - 0.047 (stat) ± 0.013 (sys)	-0.189 ± 0.054 - 0.058 (stat) ± 0.005 (sys)	-0.283 ± 0.084 - 0.095 (stat) ± 0.009 (sys)	-0.327 ± 0.076 - 0.079 (stat) ± 0.009 (sys)	0.318 ± 0.044 - 0.040 (stat) ± 0.009 (sys)	-0.141 ± 0.072 - 0.074 (stat) ± 0.005 (sys)	-0.007 ± 0.070 - 0.072 (stat) ± 0.005 (sys)	-0.004 ± 0.070 - 0.073 (stat) ± 0.006 (sys)	
15.0 - 17.0	0.349 ± 0.039 (stat) ± 0.028 (sys)	-0.142 ± 0.044 - 0.049 (stat) ± 0.007 (sys)	-0.321 ± 0.055 - 0.074 (stat) ± 0.007 (sys)	-0.316 ± 0.051 - 0.057 (stat) ± 0.008 (sys)	0.411 ± 0.041 - 0.037 (stat) ± 0.008 (sys)	0.061 ± 0.058 (stat) ± 0.005 (sys)	0.003 ± 0.061 (stat) ± 0.003 (sys)	-0.019 ± 0.054 - 0.056 (stat) ± 0.004 (sys)	
17.0 - 19.0	0.354 ± 0.049 - 0.048 (stat) ± 0.025 (sys)	-0.198 ± 0.074 - 0.084 (stat) ± 0.017 (sys)	-0.268 ± 0.063 - 0.072 (stat) ± 0.010 (sys)	-0.323 ± 0.063 - 0.072 (stat) ± 0.009 (sys)	0.305 ± 0.049 - 0.048 (stat) ± 0.013 (sys)	0.044 ± 0.073 - 0.072 (stat) ± 0.013 (sys)	0.013 ± 0.071 - 0.070 (stat) ± 0.005 (sys)	-0.094 ± 0.065 - 0.067 (stat) ± 0.004 (sys)	
Plot SelectPlot	Plot SelectPlot	Plot SelectPlot	Plot SelectPlot	Plot SelectPlot	Plot SelectPlot	Plot SelectPlot	Plot SelectPlot	Plot SelectPlot	Plot SelectPlot

Correlation tables

- ⇒ The main problem was the correlation tables...
- ⇒ We had around 80 of them...

	F_L	S_3	S_4	S_5	A_{FB}	S_7	S_8	S_9
F_L	1.00	0.06	0.00	0.03	0.04	-0.02	0.07	0.08
S_3		1.00	0.01	0.10	-0.00	-0.07	-0.01	-0.03
S_4			1.00	0.08	0.11	-0.00	0.07	0.02
S_5				1.00	0.05	-0.01	0.00	0.04
A_{FB}					1.00	0.03	-0.07	0.02
S_7						1.00	0.01	0.11
S_8							1.00	0.02
S_9								1.00

Correlation tables

⇒ The main problem was the correlation tables...

⇒ We had around 80 of them...

$q^2 = M^{*2}(\langle \text{MU} + \text{MU} \rangle)$	0.1-0.98 GeV ²								
RE	P P → B ⁰ < K [*] (892) < K ⁺ PI ⁻ > MU ⁺ MU ⁻ > X								
SQRT(S)	7000.0 GeV								
SQRT(S)	8000.0 GeV								
CORR	F_L	S_3	S_4	S_5	A_{FB}	S_7	S_8	S_9	
F_L	1.00	0.06	0.00	0.03	0.04	-0.02	0.07	0.08	
S_3	0.06	1.00	0.01	0.10	0.00	-0.07	-0.01	-0.03	
S_4	0.00	0.01	1.00	0.08	0.11	0.00	0.07	0.02	
S_5	0.03	0.10	0.08	1.00	0.05	-0.01	0.00	0.04	
A_{FB}	0.04	0.00	0.11	0.05	1.00	0.03	-0.07	0.02	
S_7	-0.02	-0.07	0.00	-0.01	0.03	1.00	0.01	0.11	
S_8	0.07	-0.01	0.07	0.00	-0.07	0.01	1.00	0.02	
S_9	0.08	-0.03	0.02	0.04	0.02	0.11	0.02	1.00	

HepData format

⇒ So HepData format is completely different then tex.

```
*dataset:  
*location: Appendix C  
*dscomment: Likelihood correlation matrix  $0.1 < q^2 < 0.98 - \{\rm GeV\}^2/c^4$ .  
*qual:  $\$q^2\$ = M^{*2}(\langle \mu^+ \mu^- \rangle)$  IN  $\text{GEV}^{*2}$ : 0.1 TO 0.98  
*reackey: P P --> B0 + X  
*obskey: CORR  
*qual: RE : P P --> B0 < K*(892) < K+ PI- > MU+ MU- > X  
*qual: Sqrt(S) IN GEV : 7000.0  
*qual: Sqrt(S) IN GEV : 8000.0  
*yheader:  $\$F_{\{\rm L\}}\$$  :  $\$S_3\$$  :  $\$S_4\$$  :  $\$S_5\$$  :  $\$A_{\{\rm FB\}}\$$  :  $\$S_7\$$  :  $\$S_8\$$  :  $\$S_9\$$   
*xheader: CORR  
*data: x : y : y : y : y : y : y : y : y  
 $\$F_{\{\rm L\}}\$$ ; 1.00; 0.06; 0.00; 0.03; 0.04; -0.02; 0.07; 0.08;  
 $\$S_3\$$ ; 0.06; 1.00; 0.01; 0.10; 0.00; -0.07; -0.01; -0.03;  
 $\$S_4\$$ ; 0.00; 0.01; 1.00; 0.08; 0.11; 0.00; 0.07; 0.02;  
 $\$S_5\$$ ; 0.03; 0.10; 0.08; 1.00; 0.05; -0.01; 0.00; 0.04;  
 $\$A_{\{\rm FB\}}\$$ ; 0.04; 0.00; 0.11; 0.05; 1.00; 0.03; -0.07; 0.02;  
 $\$S_7\$$ ; -0.02; -0.07; 0.00; -0.01; 0.03; 1.00; 0.01; 0.11;  
 $\$S_8\$$ ; 0.07; -0.01; 0.07; 0.00; -0.07; 0.01; 1.00; 0.02;  
 $\$S_9\$$ ; 0.08; -0.03; 0.02; 0.04; 0.02; 0.11; 0.02; 1.00;  
*dataend:
```


My scripts

- ⇒ Written in python.
- ⇒ Will briefly go through the some of it's functions.

```
def get_string(self):
    s = '*data: x : '
    for i in range(1, len(self.matrix_fix[0]) - 1):
        s += "y :"
    s += " y "
    self.table += s + '\n'

    for i in range(0, len(self.matrix_fix)):
        s = ''
        s += self.columns[i + 1] + " : "
        for j in range(1, len(self.matrix_fix[i])):
            #print('a',self.matrix_fix[j-1][i+1])
            s += get_ride_of_latex(self.matrix_fix[j - 1][i + 1]) + ' : '
        self.table += s + '\n'
        # self.table+= \

    print("TEST: ", self.matrix_fix)
def __unicode__(self):
    dupa=":".join(self.columns)
    dupa=dupa.replace(':', ' ', 1)
    #print(dupa)
```

My scripts

- ⇒ Written in python.
- ⇒ Will briefly go through the some of it's functions.

```
print("TEST: ", self.matrix_fix)
def __unicode__(self):
    dupa=".".join(self.columns)
    dupa=dupa.replace(':', ', ', 1)
    #print(dupa)

    return ""*dataset:
*location: Appendix G
*dscoment: %s
*yheader: %s

*data CHANGEME
%s
*dataend:
""" % (self.header, dupa, self.table)
|
class TableBuilder:
    def __init__(self):
        self.header = None
        self.columns = None
        self.matrix = []
        self.matrix_fix = []

    def add_row(self, row):
        self.matrix.append(row)

    def build(self):
        return Table(self.header, self.columns, self.matrix)

    def parse_table_header(self, line):
```


My scripts

- ⇒ Written in python.
- ⇒ Will briefly go through the some of it's functions.

```
def is_hline(self, line):  
    return line.startswith(r'\hline')  
  
def is_table_header(self, line):  
    return line.startswith(r'\subsection')  
  
def is_table_begin(self, line):  
    return line.startswith(r'\begin{tabular}')  
  
def is_table_end(self, line):  
    return line.startswith(r'\end{tabular}')  
  
def build(self):  
    self.state = ParserStateEnum.DONE  
    self.table = self.builder.build()
```

My scripts

- ⇒ Written in python.
- ⇒ Will briefly go through the some of it's functions.

```
dataset:
*location: Appendix G
*dscmment: None
*yheader:  SA_{3}$ : SA_{4}$ : SA_{5}$ : SA_{6s}$ : SA_{7}$ : SA_{8}$ : SA_{9}$

*data CHANGEME
*data: x : y :y :y :y :y :y
  SA_{3}$ : 1.00 : -0.12 : -0.18 : 0.00 : 0.01 : 0.01 : -0.05 :
  SA_{4}$ : -0.12 : 1.00 : 0.26 : -0.14 : 0.02 : -0.08 : 0.03 :
  SA_{5}$ : -0.18 : 0.26 : 1.00 : -0.13 : -0.09 : 0.02 : 0.07 :
  SA_{6s}$ : 0.00 : -0.14 : -0.13 : 1.00 : 0.0 : 0.01 : -0.01 :
  SA_{7}$ : 0.01 : 0.02 : -0.09 : 0.0 : 1.00 : 0.14 : -0.15 :
  SA_{8}$ : 0.01 : -0.08 : 0.02 : 0.01 : 0.14 : 1.00 : -0.07 :
  SA_{9}$ : -0.05 : 0.03 : 0.07 : -0.01 : -0.15 : -0.07 : 1.00 :

*dataend:
```

User case

⇒ There will be things that need to be changed for each table:

```
def get_ride_of_latex(line):  
    line2 = line  
    line2 = line2.replace(r"\\rm", r"\rm")  
    line2 = line2.replace(r"\\", "")  
    line2 = line2.replace(r'$', '')  
    line2 = line2.replace(r'\ ', '')  
    line2 = line2.replace(r'\ ', '')  
    #line2 = line2.replace(r'\gevgevcccc', 'GeV/c^4')  
    return line2
```

⇒ If you need to encode errors in HepData format:

```
*yheader: $F_{\rm L}$ : $S_{3}$ : $S_{4}$ : $S_{5}$ : $A_{\rm FB}$ : $S_{7}$ : $S_{8}$ : $S_{9}$  
*data: x : y : y : y : y : y : y : y : y  
0.10 TO 0.98 ; 0.263 +0.045, -0.044(DSYS=0.017) ; -0.036 +0.063, -0.063(DSYS=0.005) ;  
+0.058, -0.057(DSYS=0.009) ; 0.015 +0.059, -0.059(DSYS=0.006) ; 0.079 +0.076, -0.075(DSYS  
1.1 TO 2.5; 0.660 +0.083, -0.077(DSYS=0.022) ; -0.077 +0.087, -0.105(DSYS=0.005) ;  
+0.068, -0.080(DSYS=0.012) ; -0.219 +0.094, -0.104(DSYS=0.004) ; -0.098 +0.108, -0.123(DSY  
2.5 TO 4.0 ; 0.876 +0.109, -0.097(DSYS=0.017) ; 0.035 +0.098, -0.089(DSYS=0.007) ;  
+0.082, -0.090(DSYS=0.007) ; 0.068 +0.120, -0.112(DSYS=0.005) ; 0.030 +0.129, -0.131(DSYS=  
4.0 TO 6.0 ; 0.611 +0.052, -0.053(DSYS=0.017) ; 0.035 +0.069, -0.068(DSYS=0.007) ;  
+0.051, -0.052(DSYS=0.004) ; -0.016 +0.081, -0.080(DSYS=0.004) ; 0.167 +0.094, -0.091(DSYS  
5.0 TO 8.0 : 0.579 +0.046, -0.046(DSYS=0.015) : -0.042 +0.058, -0.059(DSYS=0.011) :
```

Submitting procedure

⇒ Albert and Marco have put in a procedure for submitting your results to HepData:

- You contact Alex Grecu that he prepares you a temporary slot on HepData. You will get from him number and password that you can use for logging.
- He will also prepare a JIRA task for this data submission.
- You code in your result (please remember to always have your own copy as data can be lost in HepData portal).
- Once you finish coding, your results will be reviewed/sign off by the analysis e-group.
- RD convenors are also in the loop.
- After you collected enough pokemons (:P) Alex submits this to HepData.

Summary

⇒ HepData is a tricky format that unfortunately is not similar to other ones :(

⇒ I am afraid that everyone has to adjust those scripts for they tables.

⇒ If you want to put the results to HepData do it with the paper submission!

⇒ If you already produced some results and want to copy them from tex files to Hepdata you can start with my scripts:

<http://www.physik.uzh.ch/~mchrzasz/HepData/KstarMuMu/>

⇒ Or on git:

<https://git.physik.uzh.ch/gitbucket/mchrzasz/HepData>

