





3.10pt



Searches for New Physics with LHCb





Marcin Chrząszcz

University of Zurich,
Institute of Nuclear Physics Krakow,
on behalf of LHCb collaboration

29th May 2014



University of
Zurich^{UZH}



3.10pt

LHCb detector

Lepton Flavour Violation





B decays

$$B^- \rightarrow h^+ \ell^- \ell^-$$


$$B \rightarrow \ell^+ \ell^-$$

τ decays

$$\tau^- \rightarrow \mu^- \mu^- \mu^+$$



LHCb detector



① LHCb is a forward spectrometer:


- Excellent vertex resolution.
- Efficient trigger.
- High acceptance for τ and B.
- Great Particle ID.

Lepton Flavour/Number Violation

Lepton Flavour Violation(LFV):

After μ^- was discovered it was logical to think of it as an excited e^- .

- Expected: $B(\mu \rightarrow e\gamma) \approx 10^{-4}$
- Unless another ν , in intermediate vector boson loop cancels.



I.I.Rabi:


"Who ordered that?"



- Up to this day charged LFV is being searched for in various decay modes.
- LFV was already found in neutrino sector.

Lepton Number Violation (LNV)


- Even with LFV, lepton number is a conserved quantity.
- Many new thesis predict it violation(Majorana neutrinos)
- Searched in so called Neutrinoless double β decays.




M.Chraszcz 2014

LNV in bottom decays

on-shell neutrino



virtual neutrino




- resonant production in accessible mass range
- rates depend on Majorana neutrino-lepton coupling $|V_{\mu 4}|$ (e.g. arXiv:0901.3589)
- $m_4 = m_{\ell^-, \pi^+}$

Special for B decays


Diagram without mass restriction
Cabbibo favoured for $B \rightarrow D$
Analogous to double β decay.

Virtual Majorana neutrinos

$$B^- \rightarrow D^+ \mu^- \mu^-$$



$$B^- \rightarrow D^{*+} \mu^- \mu^-$$




$\mathcal{B}(B^- \rightarrow D^+ \mu^- \mu^-) < 6.9 \times 10^{-7}$
@ 95 % CL
Based on $0.41 fb^{-1}$ 7 TeV data.

$\mathcal{B}(B^- \rightarrow D^{*+} \mu^- \mu^-) < 2.4 \times 10^{-6}$
@ 95 % CL

Phys. Rev. D85 (2012)
112004
M. Chrząszcz 2014

On-shell Majorana neutrinos


- Based on full data set $3fb^{-1}$.
- Cut based analysis.
- Normalization channel $B^+ \rightarrow J/\psi(\mu\mu)K^+$.
- Searched performed for two scenarios:
 - Short life-time neutrinos: $\tau_4 < 1ps$
 - Long life-time neutrinos: $\tau_4 \in (1, 1000)ps$



Phys. Rev. Lett.
112, 131802

M.Chrząszcz 2014

On-shell Majorana neutrinos



- In absence of signal UL. were set.
- $Br(B^- \rightarrow \pi^+ \mu^- \mu^-)$ in range 10^{-9} .
- Limits also set for the coupling $|V_{\mu 4}|^2$

$$Br(B^- \rightarrow \pi^+ \mu^- \mu^-) = \frac{G_f^4 f_B^2 f_\pi^2}{128\pi\hbar} \tau_B m_B^5 |V_{ub} V_{ud}|^2 |V_{\mu 4}|^4 \left(1 - \frac{m_4^2}{m_B^2}\right) \frac{m_4}{\Gamma_{N_4}}$$

Summary on LNV in B decays

channel	limit	
$\mathcal{B}(B^- \rightarrow \pi^+ e^- e^-)$	$< 2.3 \times 10^{-8}$	@90 % CL 
$\mathcal{B}(B^- \rightarrow K^+ e^- e^-)$	$< 3.0 \times 10^{-8}$	@90 % CL 
$\mathcal{B}(B^- \rightarrow K^{*+} e^- e^-)$	$< 2.8 \times 10^{-6}$	@90 % CL 
$\mathcal{B}(B^- \rightarrow \rho^+ e^- e^-)$	$< 2.6 \times 10^{-6}$	@90 % CL 
$\mathcal{B}(B^- \rightarrow D^+ e^- e^-)$	$< 2.6 \times 10^{-6}$	@90 % CL 
$\mathcal{B}(B^- \rightarrow D^+ e^- \mu^-)$	$< 1.8 \times 10^{-6}$	@90 % CL 
$\mathcal{B}(B^- \rightarrow \pi^+ \mu^- \mu^-)$	$< 1.3 \times 10^{-8}$	@95 % CL  <small>LHCb CERN</small>
$\mathcal{B}(B^- \rightarrow K^+ \mu^- \mu^-)$	$< 5.4 \times 10^{-7}$	@95 % CL  <small>LHCb CERN</small>
$\mathcal{B}(B^- \rightarrow D^+ \mu^- \mu^-)$	$< 6.9 \times 10^{-7}$	@95 % CL  <small>LHCb CERN</small>
$\mathcal{B}(B^- \rightarrow D^{*+} \mu^- \mu^-)$	$< 2.4 \times 10^{-6}$	@95 % CL  <small>LHCb CERN</small>
$\mathcal{B}(B^- \rightarrow D_s^+ \mu^- \mu^-)$	$< 5.8 \times 10^{-7}$	@95 % CL  <small>LHCb CERN</small>
$\mathcal{B}(B^- \rightarrow D^0 \pi^- \mu^- \mu^-)$	$< 1.5 \times 10^{-6}$	@95 % CL  <small>LHCb CERN</small>

^aBaBar, Phys. Rev. D **85**, 071103 (2012)


^bCLEO, Phys. Rev. D **65**, 111102 (2002)

^cBelle, Phys. Rev. D **84**, 071106(R), (2011)

^dLHCb, CERN-PH-EP-2012-006, arXiv:1201.5600 (2012)

$B \rightarrow \ell^+ \ell^- 1$

- A separate physics interest are LFV B decays.
- Predicted by various NP models: lepto-quarks, SUSY, GUT.
- Analysis based on 1fb^{-1} 2011 data.
- Analogues to our $B_s^0 \rightarrow \mu\mu$ analysis(Phys. Rev. Lett. 110, 021801 (2013))




- 1 Loose reselection based on topology and PID.
- 2 Classifier trained on MC signal and $b\bar{b} \rightarrow \ell\ell X$
- 3 Calibration channel: $B_{(s)}^0 \rightarrow h^+ h'^-$
- 4 Normalization Channel: $B^0 \rightarrow K^+ \pi^-$
- 5 CLs method for limit extraction.

Phys. Rev. Lett.
111, 141801 (2013)

M.Chrząszcz 2014


$B_{(s)} \rightarrow \ell^+ \ell^- 2$

- Correction to MC and DATA discrepancies.
- Excellent proxy: $B_{(s)}^0 \rightarrow hh'$.
- Fit each BDT bin for $B_{(s)}^0 \rightarrow hh'$ and extract number of events.
- Correct MC efficiency for each bin.
- Electron Bremsstrahlung corrected on $J/\psi \rightarrow ee$.




$B_{(s)} \rightarrow \ell^+ \ell^-$ Implications

- LHCb limits exceeds previous CDF by 20 times.
- CDF implications to lepto-quarks mass¹.
 - $m_{LQ}(B_s^0 \rightarrow e\mu) > 47.8(44.9) \text{ TeV}$ 90(95%) @CL.
 - $m_{LQ}(B^0 \rightarrow e\mu) > 59.3(56.3) \text{ TeV}$ 90(95%) @CL.




B decays



M.Chrząszcz 2014

$$\tau \rightarrow \mu\mu\mu$$

- 1 In SM small $\mathcal{B}(\tau^- \rightarrow \mu^-\mu^-\mu^+) \sim 10^{-50}$
- 2 NP can enhance \mathcal{B} .
- 3 Nature still hides $\tau^- \rightarrow \mu^-\mu^-\mu^+$ from us.
- 4 Current limits:



Experiment	90% CL limit
BaBar	3.3×10^{-8}
Belle	2.1×10^{-8}

- 5 Can a hadron collider change the picture?

Analysis approach

\mathcal{B} factories

LHCb, (7 TeV, 2011 data)


- 1 Clean signal: $e^+e^- \rightarrow \tau^+\tau^-$
- 2 Calculate the thrust axis
- 3 "Partial tag" the other τ
- 4 Small cross section $0.919 nb$

- 1 Inclusive τ cross section:
 $79.5 \pm 8.3 \mu b$.
- 2 $8 \times 10^{10} \tau$ produced.
- 3 Dominant contribution:
 $D_s \rightarrow \tau \nu_\tau$ (78%)
- 4 No tag possible.

Strategy

- Loose cut based selection
- Classification in 3D space:
 - invariant mass
 - decay topology (multivariate)
 - particle identification (multivariate)
- Classifier trained on simulation
- Calibration with control channel
- Normalization with $D_s \rightarrow \phi(\mu\mu)\pi$
- CLs method to extract the result

Signal & Calibration & Background channel



Signal likelihoods

particle identification


- hits in muon chambers
- energy in calorimeters
 - compatible with MIP
- RICH response

3 body decay likelihood

- vertex properties
 - vertex fit, pointing
- track quality
- isolation


Calibration

$$J/\psi \rightarrow \mu^+ \mu^-$$



Calibration

$$D_s \rightarrow \phi \pi$$





Signal likelihoods

combined signal distribution

- events distributed over 25 likelihood bins
- background estimate from mass side-bands

Signal efficiency in 3-BODY BDT vs PID BDT plane.




Extracted upper limit

LHCb
1 fb^{-1}

PLB 724
(2013) 36–45

	observed	expected	CL
$\mathcal{B}(\tau \rightarrow \mu\mu\mu)$	8.0×10^{-8}	8.3×10^{-8}	90%
	9.8×10^{-8}	10.2×10^{-8}	95%



Summary

- LFV and LNV processes are doing very well in LHCb.
- Lots of best limits already in our hands.
- Keep tune, lots of new results are coming very soon.