MC validation



Marcin Chrzaszcz Andrea Mauri Rafael Silva Coutinho





 $B^0 \to K^* \mu^- \mu^+$ meeting, CERN April 11, 2017


Intro

- ⇒ We have been ask by Vladimir to check our Filtered production.
- ⇒ So here is what we did:
- Run over the stripping on both of them.
- Truth-match: B0_BKGCAT==10
- Reweigh both Run1 and Run2 to be flat in q^2 .
- Compare ;)

Angles










- \Rightarrow Angles are oki doki.
- $\Rightarrow \cos \theta_l$ is a tinny bit getter.


RunX invariants







⇒ Overall oki-doki.












Global events







- \Rightarrow So there are differences in both kinematics and global events.
- ⇒ Naively I would expect this differences to be there.


Let's sum up what we know


- ⇒ The angular efficiencies are essentially the same.
- ⇒ Invariantes are ok.
- ⇒ All the ntupling looks oki doki.
- \Rightarrow There are differences in the kinematics and global events quantities.
- ⇒ One would naively expect them to be there.
- \Rightarrow To be sure nothing is slipping our mind I have added to comparison $B \to K^*J/\psi$ Run1 and Run2.


Reminder:


The MC $B \to K^*J/\psi$ for both Run1 and Run2 is non filtered. So if filtering is causing any problem we should see it there.


 \Rightarrow In the following plots the $B \to K^*J/\psi$ is added without any reweighing to make it flat in q^2 .














Global events








- ⇒ Everything looks ok with the filtering.
- ⇒ If you have more x-checks let me know.
- \Rightarrow Thing to think about: Move the rest of the production to LDST.

Brain storm?, credit to Andrea



Backup

