

Quark flavour anomalies of the SM



Marcin Chrzaszcz
mchrzasz@cern.ch



University of
Zurich^{UZH}









Universität Zürich,
Institute of Nuclear Physics, Polish Academy of Science

Korean Physics Society Meeting,
19 April 2017

Why electroweak penguin decays?

- In the SM allows only the charged interactions to change flavour.
 - Other interactions are flavour conserving.
- One can escape this constrain and produce $b \rightarrow s$ and $b \rightarrow d$ at loop level.
 - This kind of processes are suppressed in the SM \rightarrow Rare decays.
 - New Physics can enter in the loops.






• Operator Product Expansion and Effective Field Theory

$$H_{eff} = -\frac{4G_f}{\sqrt{2}} VV'^* \sum_i \left[\underbrace{C_i(\mu) O_i(\mu)}_{\text{left-handed}} + \underbrace{C'_i(\mu) O'_i(\mu)}_{\text{right-handed}} \right],$$



- i=1,2 Tree
- i=3-6,8 Gluon penguin
- i=7 Photon penguin
- i=9,10 EW penguin
- i=S Scalar penguin
- i=P Pseudoscalar penguin

where C_i are the Wilson coefficients and O_i are the corresponding effective operators.





- Excellent Impact Parameter (IP) resolution ($20 \mu\text{m}$).
 \Rightarrow Identify secondary vertices from heavy flavour decays
- Proper time resolution $\sim 40 - 50 \text{ fs}$.
 \Rightarrow Good separation of primary and secondary vertices.
- Excellent momentum ($\delta p/p \sim 0.5 - 1.0\%$) and inv. mass resolution.
 \Rightarrow Low combinatorial background.



- Excellent Muon identification $\epsilon_{\mu \rightarrow \mu} \sim 97\%$, $\epsilon_{\pi \rightarrow \mu} \sim 1 - 3\%$
- Good $K - \pi$ separation via RICH detectors, $\epsilon_{K \rightarrow K} \sim 95\%$,
 $\epsilon_{\pi \rightarrow K} \sim 5\%$.
 \Rightarrow Reject peaking backgrounds.
- High trigger efficiencies, low momentum thresholds.
 $B \rightarrow J/\psi X$: Trigger $\sim 90\%$.

Recent measurements of $b \rightarrow sll$

⇒ **Branching fractions:**

$$B \rightarrow K\mu^- \mu^+ \quad 1606.04731$$

$$B_s^0 \rightarrow \phi\mu^- \mu^+ \quad \text{JHEP 09 (2015) 179}$$

$$B^\pm \rightarrow \pi^\pm \mu^- \mu^+ \quad \text{JHEP 12 (2012) 125}$$

$$\Lambda_b \rightarrow \Lambda\mu^- \mu^+ \quad \text{JHEP 06 (2015) 115}$$

$$B \rightarrow \mu^- \mu^+ \quad \text{Nature 15}$$

⇒ **CP asymmetry:**

$$B^\pm \rightarrow \pi^\pm \mu^- \mu^+ \quad \text{JHEP 10 (2015) 034}$$

⇒ **Isospin asymmetry:**

$$B \rightarrow K\mu^- \mu^+ \quad \text{JHEP 06 (2014) 133}$$

⇒ **Lepton Universality:**

$$B^\pm \rightarrow K^\pm \ell \bar{\ell} \quad \text{PRL 113, (2014)}$$

⇒ **Angular:**

$$B^0 \rightarrow K^* \ell \bar{\ell} \quad \text{JHEP 02 (2016) 104}$$

$$B^{0,\pm} \rightarrow K^{*,\pm} \ell \bar{\ell} \quad \text{PRD 86 032012}$$

$$B_s^0 \rightarrow \phi \mu \mu \quad \text{JHEP 09 (2015) 179}$$

$$\Lambda_b \rightarrow \Lambda \mu^- \mu^+ \quad \text{JHEP 06 (2015) 115}$$

Recent measurements of $b \rightarrow sll$

⇒ Branching fractions:

$$B \rightarrow K\mu^- \mu^+ \quad 1606.04731$$

$$B_s^0 \rightarrow \phi\mu^- \mu^+ \quad \text{JHEP 09 (2015) 179}$$

$$B^\pm \rightarrow \pi^\pm \mu^- \mu^+ \quad \text{JHEP 12 (2012) 125}$$

$$\Lambda_b \rightarrow \Lambda\mu^- \mu^+ \quad \text{JHEP 06 (2015) 115}$$

$$B \rightarrow \mu^- \mu^+ \quad \text{Nature 15}$$

⇒ CP asymmetry:

$$B^\pm \rightarrow \pi^\pm \mu^- \mu^+ \quad \text{JHEP 10 (2015) 034}$$

⇒ Isospin asymmetry:

$$B \rightarrow K\mu^- \mu^+ \quad \text{JHEP 06 (2014) 133}$$

⇒ Lepton Universality:

$$B^\pm \rightarrow K^\pm l\bar{l} \quad \text{PRL 113, (2014)}$$

⇒ Angular:

$$B^0 \rightarrow K^* l\bar{l} \quad \text{PRD 86 032012}$$

$$B^{0,\pm} \rightarrow K^{*,\pm} l\bar{l} \quad \text{JHEP 09 (2015) 179}$$

$$B_s^0 \rightarrow \phi l\bar{l} \quad \text{JHEP 09 (2015) 179}$$

$$\Lambda_b \rightarrow \Lambda\mu^- \mu^+ \quad \text{JHEP 06 (2015) 115}$$

> 2 σ deviations from the SM

⇒ This talk is not possible to cover all flavour anomalies.

Observables in $B \rightarrow K^* \mu \mu$

⇒ The kinematics of $B^0 \rightarrow K^* \mu^- \mu^+$ decay is described by three angles θ_l , θ_k , ϕ and invariant mass of the dimuon system (q^2).

⇒ The angular distribution can be written as:

$$\frac{1}{d(\Gamma + \bar{\Gamma})/dq^2} \frac{d(\Gamma + \bar{\Gamma})}{d\cos\theta_l d\cos\theta_k d\phi} \Big|_P = \frac{9}{32\pi} \left[\frac{3}{4}(1 - F_L) \sin^2 \theta_k \right. \\ + F_L \cos^2 \theta_k + \frac{1}{4}(1 - F_L) \sin^2 \theta_k \cos 2\theta_l \\ - F_L \cos^2 \theta_k \cos 2\theta_l + S_3 \sin^2 \theta_k \sin^2 \theta_l \cos 2\phi \\ + S_4 \sin 2\theta_k \sin 2\theta_l \cos \phi + S_5 \sin 2\theta_k \sin \theta_l \cos \phi \\ + \frac{4}{3} A_{FB} \sin^2 \theta_k \cos \theta_l + S_7 \sin 2\theta_k \sin \theta_l \sin \phi \\ \left. + S_8 \sin 2\theta_k \sin 2\theta_l \sin \phi + S_9 \sin^2 \theta_k \sin^2 \theta_l \sin 2\phi \right].$$

⇒ This equation is valid in the SM for massless leptons!

Link to effective operators

⇒ The observables S_i are bilinear combinations of transversity amplitudes: $A_{\perp}^{L,R}$, $A_{\parallel}^{L,R}$, $A_0^{L,R}$.

⇒ So here is where the magic happens. At leading order the amplitudes can be written as:

$$A_{\perp}^{L,R} = \sqrt{2}Nm_B(1 - \hat{s}) \left[(C_9^{\text{eff}} + C_9^{\text{eff}'}) \mp (C_{10} + C'_{10}) + \frac{2\hat{m}_b}{\hat{s}}(C_7^{\text{eff}} + C_7^{\text{eff}'}) \right] \xi_{\perp}(E_{K^*})$$

$$A_{\parallel}^{L,R} = -\sqrt{2}Nm_B(1 - \hat{s}) \left[(C_9^{\text{eff}} - C_9^{\text{eff}'}) \mp (C_{10} - C'_{10}) + \frac{2\hat{m}_b}{\hat{s}}(C_7^{\text{eff}} - C_7^{\text{eff}'}) \right] \xi_{\perp}(E_{K^*})$$

$$A_0^{L,R} = -\frac{Nm_B(1 - \hat{s})^2}{2\hat{m}_{K^*}\sqrt{\hat{s}}} \left[(C_9^{\text{eff}} - C_9^{\text{eff}'}) \mp (C_{10} - C'_{10}) + 2\hat{m}_b(C_7^{\text{eff}} - C_7^{\text{eff}'}) \right] \xi_{\parallel}(E_{K^*}),$$

where $\hat{s} = q^2/m_B^2$, $\hat{m}_i = m_i/m_B$. The $\xi_{\parallel,\perp}$ are the soft form factors.

Link to effective operators

⇒ The observables S_i are bilinear combinations of transversity amplitudes: $A_{\perp}^{L,R}$, $A_{\parallel}^{L,R}$, $A_0^{L,R}$.

⇒ So here is where the magic happens. At leading order the amplitudes can be written as:

$$A_{\perp}^{L,R} = \sqrt{2}Nm_B(1 - \hat{s}) \left[(C_9^{\text{eff}} + C_9^{\text{eff}'}) \mp (C_{10} + C'_{10}) + \frac{2\hat{m}_b}{\hat{s}} (C_7^{\text{eff}} + C_7^{\text{eff}'}) \right] \xi_{\perp}(E_{K^*})$$

$$A_{\parallel}^{L,R} = -\sqrt{2}Nm_B(1 - \hat{s}) \left[(C_9^{\text{eff}} - C_9^{\text{eff}'}) \mp (C_{10} - C'_{10}) + \frac{2\hat{m}_b}{\hat{s}} (C_7^{\text{eff}} - C_7^{\text{eff}'}) \right] \xi_{\perp}(E_{K^*})$$


$$A_0^{L,R} = -\frac{Nm_B(1 - \hat{s})^2}{2\hat{m}_{K^*}\sqrt{\hat{s}}} \left[(C_9^{\text{eff}} - C_9^{\text{eff}'}) \mp (C_{10} - C'_{10}) + 2\hat{m}_b(C_7^{\text{eff}} - C_7^{\text{eff}'}) \right] \xi_{\parallel}(E_{K^*}),$$

where $\hat{s} = q^2/m_B^2$, $\hat{m}_i = m_i/m_B$. The $\xi_{\parallel,\perp}$ are the soft form factors.

⇒ Now we can construct observables that cancel the ξ soft form factors at leading order:

$$P'_5 = \frac{S_5 + \bar{S}_5}{2\sqrt{F_L(1 - F_L)}}$$

- Signal modelled by a sum of two Crystal-Ball functions.
 - Shape is defined using $B \rightarrow J/\psi K^*$ and corrected for q^2 dependency.
 - Combinatorial background modelled by exponent.
- $K\pi$ system:
 - Beside the K^* resonance there might be a tail from other higher mass states.
 - We modelled it in the analysis.
 - Reduced the systematic compared to previous analysis.
- In total we found 2398 ± 57 candidates in the $(0.1, 19) \text{ GeV}^2/c^4$ q^2 region.
 - 624 ± 30 candidates in the theoretically the most interesting $(1.1 - 6.0) \text{ GeV}^2/c^4$ region.





- Detector distorts our angular distribution.
- We need to model this effect.
- Full 4D function is used:

$$\epsilon(\cos \theta_l, \cos \theta_k, \phi, q^2) =$$






$$\sum_{ijkl} c_{ijkl} P_i(\cos \theta_l) P_j(\cos \theta_k) P_k(\phi) P_l(q^2),$$





where P_i is the Legendre polynomial of order i .





- We use up to 4th, 5th, 6th, 5th order for the $\cos \theta_l, \cos \theta_k, \phi, q^2$.
- 600 terms in total!




- We tested our unfolding procedure on $B \rightarrow J/\psi K^*$.
- The result is in perfect agreement with other experiments and our different analysis of this decay.
















- Tension gets confirmed!
- The two bins deviate by 2.8 and 3.0 σ from the SM prediction.
- Result compatible with previous result.



- Suppressed by $\frac{f_s}{f_d}$.
- Cleaner because of narrow ϕ resonance.
- 3.3σ deviation in the SM in the $1 - 6 \text{ GeV}^2/c^4$ bin.
- Angular part in agreement with the SM (S_5 is not accessible).






- In total ~ 300 candidates in data set.
- Decay not present in the low q^2 .
- For the bins in which we have $> 3 \sigma$ significance the forward backward asymmetry is measured for the hadronic and leptonic system.





- Challenging analysis due to bremsstrahlung.
- Migration of events modeled by MC.
- Correct for bremsstrahlung.
- Take double ratio with $B^+ \rightarrow J/\psi K^+$ to cancel systematics.
- In 3fb^{-1} , LHCb measures $R_K = 0.745^{+0.090}_{-0.074}(\text{stat.})^{+0.036}_{-0.036}(\text{syst.})$
- Consistent with the SM at 2.6σ .

$$R_K = \frac{\int_{q^2=1 \text{ GeV}^2/c^4}^{q^2=6 \text{ GeV}^2/c^4} (dB[B^+ \rightarrow K^+ \mu^+ \mu^-]/dq^2) dq^2}{\int_{q^2=1 \text{ GeV}^2/c^4}^{q^2=6 \text{ GeV}^2/c^4} (dB[B^+ \rightarrow K^+ e^+ e^-]/dq^2) dq^2} = 1 \pm \mathcal{O}(10^{-3})$$




- A fit prepared by S. Descotes-Genon, L. Hofer, J. Matias, J. Virto.
- The data can be explained by modifying the C_9 Wilson coefficient.
- Overall there is $> 4 \sigma$ discrepancy wrt. the SM prediction.




If not NP?

- We are not there yet!
- There might be something not taken into account in the theory.
- Resonances (J/ψ , $\psi(2S)$) tails can mimic NP effects.
- There might be some non factorizable QCD corrections.
" However, the central value of this effect would have to be significantly larger than expected on the basis of existing estimates" D. Straub, [arXiv:1503.06199](https://arxiv.org/abs/1503.06199) .



If not NP?

- We are not there yet!
- There might be something not taken into account in the theory.
- Resonances (J/ψ , $\psi(2S)$) tails can mimic NP effects.
- There might be some non factorizable QCD corrections.
” However, the central value of this effect would have to be significantly larger than expected on the basis of existing estimates” D. Straub, [arXiv:1503.06199](https://arxiv.org/abs/1503.06199) .



Grab it While it's Hot!

- ⇒ Yesterday(18.04) we shown a new preliminary result: [CERN Seminar](#)
- ⇒ We measured the ratio:

$$R_{K^*} = \frac{\mathcal{B}(B \rightarrow K^* \mu \mu)}{\mathcal{B}(B \rightarrow K^* e e)}$$



Grab it While it's Hot!

- ⇒ Yesterday(18.04) we shown a new preliminary result: [CERN Seminar](#)
- ⇒ We measured the ratio:

$$R_{K^*} = \frac{\mathcal{B}(B \rightarrow K^* \mu \mu)}{\mathcal{B}(B \rightarrow K^* e e)}$$

⇒ Measurement performed in two q^2 bins.



⇒ Normalized in double ratio to $B \rightarrow K^* J/\psi$.



Grab it While it's Hotter!


⇒ Today(19.04) there was already first paper with the phenomenological work about this measurement: arxiv::1704.05340 J. Matias, et. al.

1D Hyp.	All					LFUV				
	Best fit	1σ	2σ	Pull _{SM}	p-value	Best fit	1σ	2σ	Pull _{SM}	p-value
$C_{9\mu}^{\text{NP}}$	-1.10	[-1.27, -0.92]	[-1.43, -0.74]	5.7	72	-1.76	[-2.36, -1.23]	[-3.04, -0.76]	3.9	69
$C_{9\mu}^{\text{NP}} = -C_{10\mu}^{\text{NP}}$	-0.61	[-0.73, -0.48]	[-0.87, -0.36]	5.2	61	-0.66	[-0.84, -0.48]	[-1.04, -0.32]	4.1	78
$C_{9\mu}^{\text{NP}} = -C'_{9\mu}$	-1.01	[-1.18, -0.84]	[-1.33, -0.65]	5.4	66	-1.64	[-2.12, -1.05]	[-2.52, -0.49]	3.2	31
$C_{9\mu}^{\text{NP}} = -3C_{9e}^{\text{NP}}$	-1.06	[-1.23, -0.89]	[-1.39, -0.71]	5.8	74	-1.35	[-1.82, -0.95]	[-2.38, -0.59]	4.0	71



There is more!

- There is one other Lepton Universality Violation decay recently measured by LHCb.
- $R(D^*) = \frac{\mathcal{B}(B \rightarrow D^* \tau \nu)}{\mathcal{B}(B \rightarrow D^* \mu \nu)}$
- Clean SM prediction: $R(D^*) = 0.252(3)$, [PRD 85 094025 \(2012\)](#)
- LHCb result: $R(D^*) = 0.336 \pm 0.027 \pm 0.030$
- HFAG average: $R(D^*) = 0.322 \pm 0.022$
- 4.0σ discrepancy wrt. SM.



Conclusions

- Clear tensions wrt. SM predictions!
- Measurements cluster in the same direction.
- We are not opening the champagne yet!
- Still need improvement both on theory and experimental side.
- More data will shade a light on the matter.
- Time will tell if this is QCD+fluctuations or new Physics:

Conclusions

- Clear tensions wrt. SM predictions!
- Measurements cluster in the same direction.
- We are not opening the champagne yet!
- Still need improvement both on theory and experimental side.
- More data will shade a light on the matter.
- Time will tell if this is QCD+fluctuations or new Physics:

”... when you have eliminated all the Standard Model explanations, whatever remains, however improbable, must be New Physics.”

Prof. Joaquim Matias

Backup



Theory implications

Coefficient	Best fit	1σ	3σ	Pull _{SM}	p-value (%)
C_7^{NP}	-0.02	[-0.04, -0.00]	[-0.07, 0.04]	1.1	16.0
C_9^{NP}	-1.11	[-1.32, -0.89]	[-1.71, -0.40]	4.5	62.0
C_{10}^{NP}	0.58	[0.34, 0.84]	[-0.11, 1.41]	2.5	25.0
$C_{7'}^{\text{NP}}$	0.02	[-0.01, 0.04]	[-0.05, 0.09]	0.7	15.0
$C_{9'}^{\text{NP}}$	0.49	[0.21, 0.77]	[-0.33, 1.35]	1.8	19.0
$C_{10'}^{\text{NP}}$	-0.27	[-0.46, -0.08]	[-0.84, 0.28]	1.4	17.0
$C_9^{\text{NP}} = C_{10}^{\text{NP}}$	-0.21	[-0.40, 0.00]	[-0.74, 0.55]	1.0	16.0
$C_9^{\text{NP}} = -C_{10}^{\text{NP}}$	-0.69	[-0.88, -0.51]	[-1.27, -0.18]	4.1	55.0
$C_{9'}^{\text{NP}} = C_{10'}^{\text{NP}}$	-0.09	[-0.35, 0.17]	[-0.88, 0.66]	0.3	14.0
$C_{9'}^{\text{NP}} = -C_{10'}^{\text{NP}}$	0.20	[0.08, 0.32]	[-0.15, 0.56]	1.7	19.0
$C_9^{\text{NP}} = -C_{9'}^{\text{NP}}$	-1.09	[-1.28, -0.88]	[-1.62, -0.42]	4.8	72.0
$C_9^{\text{NP}} = -C_{10}^{\text{NP}}$ $= -C_{9'}^{\text{NP}} = -C_{10'}^{\text{NP}}$	-0.68	[-0.49, -0.49]	[-1.36, -0.15]	3.9	50.0
$C_9^{\text{NP}} = -C_{10}^{\text{NP}}$ $= C_{9'}^{\text{NP}} = -C_{10'}^{\text{NP}}$	-0.17	[-0.29, -0.06]	[-0.54, 0.18]	1.5	18.0

Table 2: *Best-fit points, confidence intervals, pulls for the SM hypothesis and p-values for different one-dimensional NP scenarios.*

If not NP?

- How about our clean P_i observables?
- The QCD cancel as mentioned only at leading order.
- Comparison to normal observables with the optimised ones.



Transversity amplitudes

⇒ One can link the angular observables to transversity amplitudes

$$J_{1s} = \frac{(2 + \beta_\ell^2)}{4} \left[|A_\perp^L|^2 + |A_\parallel^L|^2 + |A_\perp^R|^2 + |A_\parallel^R|^2 \right] + \frac{4m_\ell^2}{q} \operatorname{Re} \left(A_\perp^L A_\perp^{R*} + A_\parallel^L A_\parallel^{R*} \right),$$

$$J_{1c} = |A_0^L|^2 + |A_0^R|^2 + \frac{4m_\ell^2}{q} \left[|A_t|^2 + 2\operatorname{Re}(A_0^L A_0^{R*}) \right] + \beta_\ell^2 |A_S|^2,$$

$$J_{2s} = \frac{\beta_\ell^2}{4} \left[|A_\perp^L|^2 + |A_\parallel^L|^2 + |A_\perp^R|^2 + |A_\parallel^R|^2 \right], \quad J_{2c} = -\beta_\ell^2 \left[|A_0^L|^2 + |A_0^R|^2 \right],$$

$$J_3 = \frac{1}{2} \beta_\ell^2 \left[|A_\perp^L|^2 - |A_\parallel^L|^2 + |A_\perp^R|^2 - |A_\parallel^R|^2 \right], \quad J_4 = \frac{1}{\sqrt{2}} \beta_\ell^2 \left[\operatorname{Re}(A_0^L A_\parallel^{L*} + A_0^R A_\parallel^{R*}) \right],$$

$$J_5 = \sqrt{2} \beta_\ell \left[\operatorname{Re}(A_0^L A_\perp^{L*} - A_0^R A_\perp^{R*}) - \frac{m_\ell}{\sqrt{q^2}} \operatorname{Re}(A_\parallel^L A_S^* + A_\parallel^{R*} A_S) \right],$$

$$J_{6s} = 2\beta_\ell \left[\operatorname{Re}(A_\parallel^L A_\perp^{L*} - A_\parallel^R A_\perp^{R*}) \right], \quad J_{6c} = 4\beta_\ell \frac{m_\ell}{\sqrt{q^2}} \operatorname{Re}(A_0^L A_S^* + A_0^{R*} A_S),$$

$$J_7 = \sqrt{2} \beta_\ell \left[\operatorname{Im}(A_0^L A_\parallel^{L*} - A_0^R A_\parallel^{R*}) + \frac{m_\ell}{\sqrt{q^2}} \operatorname{Im}(A_\perp^L A_S^* - A_\perp^{R*} A_S) \right],$$

$$J_8 = \frac{1}{\sqrt{2}} \beta_\ell^2 \left[\operatorname{Im}(A_0^L A_\perp^{L*} + A_0^R A_\perp^{R*}) \right], \quad J_9 = \beta_\ell^2 \left[\operatorname{Im}(A_\parallel^{L*} A_\perp^L + A_\parallel^{R*} A_\perp^R) \right],$$

Link to effective operators

⇒ So here is where the magic happens. At leading order the amplitudes can be written as (soft form factors):

$$A_{\perp}^{L,R} = \sqrt{2} N m_B (1 - \hat{s}) \left[(C_9^{\text{eff}} + C_9^{\text{eff}'}) \mp (C_{10} + C'_{10}) + \frac{2\hat{m}_b}{\hat{s}} (C_7^{\text{eff}} + C_7^{\text{eff}'}) \right] \xi_{\perp}(E_{K^*})$$

$$A_{\parallel}^{L,R} = -\sqrt{2} N m_B (1 - \hat{s}) \left[(C_9^{\text{eff}} - C_9^{\text{eff}'}) \mp (C_{10} - C'_{10}) + \frac{2\hat{m}_b}{\hat{s}} (C_7^{\text{eff}} - C_7^{\text{eff}'}) \right] \xi_{\parallel}(E_{K^*})$$

$$A_0^{L,R} = -\frac{N m_B (1 - \hat{s})^2}{2\hat{m}_{K^*} \sqrt{\hat{s}}} \left[(C_9^{\text{eff}} - C_9^{\text{eff}'}) \mp (C_{10} - C'_{10}) + 2\hat{m}_b (C_7^{\text{eff}} - C_7^{\text{eff}'}) \right] \xi_{\parallel}(E_{K^*}),$$

where $\hat{s} = q^2/m_B^2$, $\hat{m}_i = m_i/m_B$. The $\xi_{\parallel,\perp}$ are the form factors.

Link to effective operators

⇒ So here is where the magic happens. At leading order the amplitudes can be written as (soft form factors):

$$A_{\perp}^{L,R} = \sqrt{2} N m_B (1 - \hat{s}) \left[(C_9^{\text{eff}} + C_9^{\text{eff}'}) \mp (C_{10} + C'_{10}) + \frac{2\hat{m}_b}{\hat{s}} (C_7^{\text{eff}} + C_7^{\text{eff}'}) \right] \xi_{\perp}(E_{K^*})$$

$$A_{\parallel}^{L,R} = -\sqrt{2} N m_B (1 - \hat{s}) \left[(C_9^{\text{eff}} - C_9^{\text{eff}'}) \mp (C_{10} - C'_{10}) + \frac{2\hat{m}_b}{\hat{s}} (C_7^{\text{eff}} - C_7^{\text{eff}'}) \right] \xi_{\perp}(E_{K^*})$$

$$A_0^{L,R} = -\frac{N m_B (1 - \hat{s})^2}{2\hat{m}_{K^*} \sqrt{\hat{s}}} \left[(C_9^{\text{eff}} - C_9^{\text{eff}'}) \mp (C_{10} - C'_{10}) + 2\hat{m}_b (C_7^{\text{eff}} - C_7^{\text{eff}'}) \right] \xi_{\parallel}(E_{K^*}),$$

where $\hat{s} = q^2/m_B^2$, $\hat{m}_i = m_i/m_B$. The $\xi_{\parallel,\perp}$ are the form factors.

⇒ Now we can construct observables that cancel the ξ form factors at leading order:

$$P'_5 = \frac{J_5 + \bar{J}_5}{2\sqrt{-(J_2^c + \bar{J}_2^c)(J_2^s + \bar{J}_2^s)}}$$


$B^0 \rightarrow K^* \mu^- \mu^+$ kinematics

\Rightarrow The kinematics of $B^0 \rightarrow K^* \mu^- \mu^+$ decay is described by three angles θ_l , θ_k , ϕ and invariant mass of the dimuon system (q^2).


$\Rightarrow \cos \theta_k$: the angle between the direction of the kaon in the K^* (\bar{K}^*) rest frame and the direction of the K^* (\bar{K}^*) in the B^0 (\bar{B}^0) rest frame.

$\Rightarrow \cos \theta_l$: the angle between the direction of the μ^- (μ^+) in the dimuon rest frame and the direction of the dimuon in the B^0 (\bar{B}^0) rest frame.


$\Rightarrow \phi$: the angle between the plane containing the μ^- and μ^+ and the plane containing the kaon and pion from the K^* .



(a) θ_K and θ_l definitions for the B^0 decay



(b) ϕ definition for the B^0 decay



(c) ϕ definition for the \bar{B}^0 decay

$B^0 \rightarrow K^* \mu^- \mu^+$ kinematics

\Rightarrow The kinematics of $B^0 \rightarrow K^* \mu^- \mu^+$ decay is described by three angles θ_l, θ_k, ϕ and invariant mass of the dimuon system (q^2).

$$\begin{aligned} \frac{d^4\Gamma}{dq^2 d\cos\theta_K d\cos\theta_l d\phi} = & \frac{9}{32\pi} \left[J_{1s} \sin^2\theta_K + J_{1c} \cos^2\theta_K + (J_{2s} \sin^2\theta_K + J_{2c} \cos^2\theta_K) \cos 2\theta_l \right. \\ & + J_3 \sin^2\theta_K \sin^2\theta_l \cos 2\phi + J_4 \sin 2\theta_K \sin 2\theta_l \cos \phi + J_5 \sin 2\theta_K \sin \theta_l \cos \phi \\ & + (J_{6s} \sin^2\theta_K + J_{6c} \cos^2\theta_K) \cos \theta_l + J_7 \sin 2\theta_K \sin \theta_l \sin \phi + J_8 \sin 2\theta_K \sin 2\theta_l \sin \phi \\ & \left. + J_9 \sin^2\theta_K \sin^2\theta_l \sin 2\phi \right], \end{aligned}$$

\Rightarrow This is the most general expression of this kind of decay.

\Rightarrow The CP averaged angular observables are defined:

$$S_i = \frac{J_i + \bar{J}_i}{(d\Gamma + d\bar{\Gamma})/dq^2}$$

Link to effective operators

⇒ The observables J_i are bilinear combinations of transversity amplitudes: $A_{\perp}^{L,R}$, $A_{\parallel}^{L,R}$, $A_0^{L,R}$.

⇒ So here is where the magic happens. At leading order the amplitudes can be written as:

$$A_{\perp}^{L,R} = \sqrt{2}Nm_B(1 - \hat{s}) \left[(C_9^{\text{eff}} + C_9^{\text{eff}'}) \mp (C_{10} + C'_{10}) + \frac{2\hat{m}_b}{\hat{s}} (C_7^{\text{eff}} + C_7^{\text{eff}'}) \right] \xi_{\perp}(E_{K^*})$$

$$A_{\parallel}^{L,R} = -\sqrt{2}Nm_B(1 - \hat{s}) \left[(C_9^{\text{eff}} - C_9^{\text{eff}'}) \mp (C_{10} - C'_{10}) + \frac{2\hat{m}_b}{\hat{s}} (C_7^{\text{eff}} - C_7^{\text{eff}'}) \right] \xi_{\perp}(E_{K^*})$$

$$A_0^{L,R} = -\frac{Nm_B(1 - \hat{s})^2}{2\hat{m}_{K^*}\sqrt{\hat{s}}} \left[(C_9^{\text{eff}} - C_9^{\text{eff}'}) \mp (C_{10} - C'_{10}) + 2\hat{m}_b(C_7^{\text{eff}} - C_7^{\text{eff}'}) \right] \xi_{\parallel}(E_{K^*}),$$

where $\hat{s} = q^2/m_B^2$, $\hat{m}_i = m_i/m_B$. The $\xi_{\parallel,\perp}$ are the soft form factors.

Link to effective operators

⇒ The observables J_i are bilinear combinations of transversity amplitudes: $A_{\perp}^{L,R}$, $A_{\parallel}^{L,R}$, $A_0^{L,R}$.

⇒ So here is where the magic happens. At leading order the amplitudes can be written as:

$$A_{\perp}^{L,R} = \sqrt{2} N m_B (1 - \hat{s}) \left[(c_9^{\text{eff}} + c_9^{\text{eff}'}) \mp (c_{10} + c'_{10}) + \frac{2\hat{m}_b}{\hat{s}} (c_7^{\text{eff}} + c_7^{\text{eff}'}) \right] \xi_{\perp}(E_{K^*})$$

$$A_{\parallel}^{L,R} = -\sqrt{2} N m_B (1 - \hat{s}) \left[(c_9^{\text{eff}} - c_9^{\text{eff}'}) \mp (c_{10} - c'_{10}) + \frac{2\hat{m}_b}{\hat{s}} (c_7^{\text{eff}} - c_7^{\text{eff}'}) \right] \xi_{\perp}(E_{K^*})$$

$$A_0^{L,R} = -\frac{N m_B (1 - \hat{s})^2}{2\hat{m}_{K^*} \sqrt{\hat{s}}} \left[(c_9^{\text{eff}} - c_9^{\text{eff}'}) \mp (c_{10} - c'_{10}) + 2\hat{m}_b (c_7^{\text{eff}} - c_7^{\text{eff}'}) \right] \xi_{\parallel}(E_{K^*}),$$

where $\hat{s} = q^2/m_B^2$, $\hat{m}_i = m_i/m_B$. The $\xi_{\parallel,\perp}$ are the soft form factors.

⇒ Now we can construct observables that cancel the ξ soft form factors at leading order:

$$P'_5 = \frac{J_5 + \bar{J}_5}{2\sqrt{-(J_2^c + \bar{J}_2^c)(J_2^s + \bar{J}_2^s)}}$$

Symmetries in $B \rightarrow K^* \mu \mu$

⇒ We have 12 angular coefficients (S_i).

⇒ There exist 4 symmetry transformations that leave the angular distributions unchanged:


$$n_{\parallel} = \begin{pmatrix} A_{\parallel}^L \\ A_{R^*}^L \end{pmatrix}, \quad n_{\perp} = \begin{pmatrix} A_{\perp}^L \\ -A_{\perp}^L \end{pmatrix}, \quad n_0 = \begin{pmatrix} A_0^L \\ A_0^L \end{pmatrix}.$$

$$n_i' = U n_i = \begin{bmatrix} e^{i\phi_L} & 0 \\ 0 & e^{-i\phi_R} \end{bmatrix} \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} \cosh i\tilde{\theta} & -\sinh i\tilde{\theta} \\ -\sinh i\tilde{\theta} & \cosh i\tilde{\theta} \end{bmatrix} n_i.$$

⇒ Using this symmetries one can show that there are 8 independent observables. The pdf can be written as:

$$\frac{1}{d(\Gamma + \bar{\Gamma})/dq^2} \frac{d(\Gamma + \bar{\Gamma})}{d\cos\theta_l d\cos\theta_k d\phi} \Big|_P = \frac{9}{32\pi} \left[\frac{3}{4}(1 - F_L) \sin^2 \theta_k \right. \\ \left. + F_L \cos^2 \theta_k + \frac{1}{4}(1 - F_L) \sin^2 \theta_k \cos 2\theta_l \right. \\ \left. - F_L \cos^2 \theta_k \cos 2\theta_l + S_3 \sin^2 \theta_k \sin^2 \theta_l \cos 2\phi \right. \\ \left. + S_4 \sin 2\theta_k \sin 2\theta_l \cos \phi + S_5 \sin 2\theta_k \sin \theta_l \cos \phi \right. \\ \left. + \frac{4}{3} A_{\text{FB}} \sin^2 \theta_k \cos \theta_l + S_7 \sin 2\theta_k \sin \theta_l \sin \phi \right. \\ \left. + S_8 \sin 2\theta_k \sin 2\theta_l \sin \phi + S_9 \sin^2 \theta_k \sin^2 \theta_l \sin 2\phi \right].$$

- Thanks to Method of Moments there was the possibility to measure a new observable S_{6c} .



- Measurement is consistent with the SM prediction.

- With the full data set (3fb^{-1}) we performed angular analysis in $0.0004 < q^2 < 1 \text{ GeV}^2/c^4$.
- Electrons channels are extremely challenging experimentally:
 - Bremsstrahlung.
 - Trigger efficiencies.
- Determine the angular observables: F_L , $A_T^{(2)}$, A_T^{Re} , A_T^{Im} :
- Results in full agreement with the SM.
- Similar strength on C_7 Wilson coefficient as from $b \rightarrow s\gamma$ decays.

