MVA selection for $au o 3\mu$


Marcin Chrząszcz^{1,2}

¹ University of Zurich, ² Institute of Nuclear Physics, Krakow,

January 6, 2014








Blending - THE END

Binning optimisation

 η background

Plan




Where did we end last year

- We decided to use Blending technique with MatrixNet.
- Jon updated the mixing numbers → Need to retrain everything and hope things are status quo.
- People thanks to $B^0_s \to \mu\mu$ will ask about mass correlation \to Need to check to shut them up. Big big surprise.

M.Chrzaszcz 2013 3/14

Status quo

- I was to lazy to retrain all TMVA stuff.
- So just compared: GeoMVA(we have it for free), MN standalone, Blending.
- Again Blending wins.




M.Chrząszcz 2013

Update on analysis Blending - THE END 4 / 14

Status quo

- I was to lazy to retrain all TMVA stuff
- So just compared: GeoMVA(we have it for free), MN standalone, Blending.
- Again Blending wins. \rightarrow I am happy.




M.Chrząszcz 2013

Update on analysis Blending - THE END 5 / 14

Status quo


- For sanity check I also have looked into 2011 data.
- Result stays the same.





Update on analysis Blending - THE END 6 / 14

How does MN look like.

- So MN is Gradient BDT. (name in the ntuple: MN_BLEND)
- Applied transformation to make flat signal distribution. (name in the ntuple:
 MN BLEND FLAT).








Mass correlation - Sit down, this is hardcore

- So lets check the correlation for my blending using MN.
- Correlation is flat and arround 3.5%.
 Everything seems to be ok.
- Now the scary part. I checked this for GeoMVA.




0.00

-0.05

-0.10

Mass correlation - Sit down, this is hardcore

- I saw 30% correlation in our previous GEOMVA!
- Before I got a heart attack I had thought: Making clarifies flat has to change the correlation value. So GeoMVA was transformed to be flat, for 1:1 comaprison.
- Never the less 15% in GeoMVA remains;




MVA conclusions

- Blending is the best performing MVA.
- Mass correlation is reasonable low.
- GeoMVa mystery: Different RECO might have something to do with what we saw. It also could explain why GeoMVA is so efficient now(correlation might make the bck level artificially lower)
 ← this idea was after midnight and 3 pines of beer so I might now change my mind.

Update on analysis Blending - THE END 10 / 14

Binning


- Performed binning optimisation(with old trigger strategy and old classifier(before Jons updates)).
- All this will be updated of coz with Pauls new results + new clarifies
- For 2012 for binning with eta veto: PID: 0 0.10 0.42 0.75 0.81 0.86 1.00 BLEND FLAT: 0. 0.05 0.38 0.63 0.71 0.82 1.00



M.Chrzaszcz 2013

Binning - details

- Binning uses CLs method, not enough time to play with other FOM.
- Optimisation requires at least 8 events in the bin.
- Minimum bin size is 0.06.
- The sieve is 0.01



M.Chrząszcz 2013

Binning - details

- Did all the fits to η bck.
- · Fits are far more stable then last year.
- Will try to make a single fit to all bins to reduce the systematics.
- All fits are here: LINK
- I was lazy and didn't want to latex thing that will change...
- The same with fits with η veto....
- MC/Data comparison doing Splot: LINK

 M.Chrząszcz 2013

 Update on analysis
 η background
 13 / 14

The Plan

- 1 If we want to move with this for Moriond we would need to have a limit soon.
- I would propose the attemnt to have on next week a preliminary limit.
- What we need:
- $D_s \rightarrow \phi \pi$. Done for 2011 and 2012. Will appear today on my page.
- New binning: 6 hours of computing one scenario
 → tmr will have it.
- Fits(with η veto: Wed).
- Calibration: Paul?
- Limit(lets aim 2011+2012). I have script from Diego with the combination for $B_s^0 \to \mu\mu$.
- But it is never easy to run those things(2-3 days).
- Lets try to make an effort?

M.Chrzaszcz 2013 14 / 14