

Searches for Lepton Flavour Violation and Lepton Number Violation in Hadron Decays

Paul Seyfert

on behalf of the LHCb collaboration
including results from the B factories

Heidelberg University

Flavour Physics and CP Violation 2012

SPONSORED BY THE





Outline

1 Lepton Number Violation

- $D^- \rightarrow \pi^+ / K^+ \ell^- \ell^-$
- $B^- \rightarrow h^+ \ell^- \ell^-$
- $B^- \rightarrow D^0 \pi^+ \mu^- \mu^-$

2 Lepton Flavour Violation

- $D^- \rightarrow h^- \ell^+ \ell'^-$
- $B^- \rightarrow \pi^- / K^- \tau^\pm \ell^\mp$

3 $\tau^- \rightarrow \mu^- \mu^- \mu^+$ at hadron colliders – first time shown!


Lepton Number Violation

- $D^- \rightarrow \pi^+ / K^+ \ell^- \ell^-$
- $B^- \rightarrow h^+ \ell^- \ell^-$
- $B^- \rightarrow D^0 \pi^+ \mu^- \mu^-$



Lepton Number Violation


- Lepton number is conserved in the Standard Model
- Lepton number violation introduced in many New Physics models, e. g.:
 - 4th quark generation
([arXiv:hep-ph/1106.0343](https://arxiv.org/abs/hep-ph/1106.0343))
 - SO(10) SUSY GUT
([arXiv:hep-ph/9501298](https://arxiv.org/abs/hep-ph/9501298))
 - exotic Higgs
(A. Zee, *Phys. Lett.* B93, 389 (1980))
- Majorana neutrinos violate Lepton Number
classic search: neutrinoless double beta-decay
(→ [Dongming Mei's talk](#))
- Similar fundamental process with mesons





LNV in meson decays

$$D^+ \rightarrow \pi^- \mu^+ \mu^+$$




- resonant production in accessible mass range
- rates depend on Majorana neutrino-lepton coupling $|V_{\mu 4}|$
(e.g. [arXiv:0901.3589](https://arxiv.org/abs/0901.3589))
- $m_4 = m_{\ell^-, \pi^+}$






Lepton Number Violation in meson decays

$$D^+ \rightarrow \pi^- \mu^+ \mu^+$$



- resonant production in accessible mass range
- rates depend on Majorana neutrino-lepton coupling $|V_{\mu 4}|$
(e.g. [arXiv:0901.3589](https://arxiv.org/abs/0901.3589))
- $m_4 = m_{\ell^-, \pi^+}$



Status of 2009

[arXiv:0901.3589](https://arxiv.org/abs/0901.3589)





limits on LNV in charm decays

charm decays

- latest limits from BaBar
- includes Lepton Number and Flavour Violation
- comprehensive list of D^+ , D_s^+ , and Λ_c^+ decays



Decay mode	Yield (events)	Eff. (%)	90% CL (10^{-4})	BR UL 90% CL (10^{-6})	BF UL (10^{-6})
$D^+ \rightarrow \pi^- e^+ e^+$	$4.7 \pm 4.7 \pm 0.5$	3.16	6.8	1.9	
$D^+ \rightarrow \pi^- \mu^+ \mu^+$	$-3.1 \pm 1.2 \pm 0.5$	0.70	7.5	2.0	
$D^+ \rightarrow \pi^- \mu^+ e^+$	$-5.1 \pm 4.2 \pm 2.0$	1.72	7.4	2.0	
$D_s^+ \rightarrow \pi^- e^+ e^+$	$-5.7 \pm 14. \pm 3.4$	6.84	1.8	4.1	
$D_s^+ \rightarrow \pi^- \mu^+ \mu^+$	$0.6 \pm 5.1 \pm 2.7$	1.05	6.2	14	
$D_s^+ \rightarrow \pi^- \mu^+ e^+$	$-0.2 \pm 7.9 \pm 0.6$	2.23	3.6	8.4	
$D^+ \rightarrow K^- e^+ e^+$	$-2.8 \pm 2.4 \pm 0.2$	2.67	3.1	0.9	
$D^+ \rightarrow K^- \mu^+ \mu^+$	$7.2 \pm 5.4 \pm 1.6$	0.80	37	10	
$D^+ \rightarrow K^- \mu^+ e^+$	$-11.6 \pm 4.0 \pm 3.1$	1.52	6.8	1.9	
$D_s^+ \rightarrow K^- e^+ e^+$	$2.3 \pm 7.9 \pm 3.3$	4.10	2.1	5.2	
$D_s^+ \rightarrow K^- \mu^+ \mu^+$	$-2.3 \pm 5.0 \pm 2.8$	0.98	5.3	13	
$D_s^+ \rightarrow K^- \mu^+ e^+$	$-14.0 \pm 8.4 \pm 2.0$	2.26	2.4	6.1	
$\Lambda_c^+ \rightarrow \bar{p} e^+ e^+$	$-1.5 \pm 4.2 \pm 1.5$	5.14	0.4	2.7	
$\Lambda_c^+ \rightarrow \bar{p} \mu^+ \mu^+$	$-0.0 \pm 2.1 \pm 0.6$	0.94	1.4	9.4	
$\Lambda_c^+ \rightarrow \bar{p} \mu^+ e^+$	$10.1 \pm 5.8 \pm 3.5$	2.50	2.3	16	




Phys. Rev. D 84, 072006 (2011)



limits on LNV in charm decays

charm decays

- latest limits from BaBar
- includes Lepton Number and Flavour Violation
- comprehensive list of D^+ , D_s^+ , and Λ_c^+ decays
- best limit for most channels





Lepton Number Violation in bottom decays


$$B^- \rightarrow h^+ \ell^- \ell^-$$






Lepton Number Violation in bottom decays

on-shell neutrino



virtual neutrino



- resonant production in accessible mass range
- rates depend on Majorana neutrino-lepton coupling $|V_{\mu 4}|$ (e.g. [arXiv:0901.3589](https://arxiv.org/abs/0901.3589))
- $m_4 = m_{\ell^-, \pi^+}$

special for B decays

diagram without mass restriction
Cabbibo favoured for $B \rightarrow D$





search for Majorana neutrinos at LHCb

- searches at B factories in Liang Sun's talk
- LHCb: search for

- $B^- \rightarrow \pi^+ \mu^- \mu^-$
- $B^- \rightarrow D^+ \mu^- \mu^-$
- $B^- \rightarrow D_s^+ \mu^- \mu^-$
- $B^- \rightarrow D^{*+} \mu^- \mu^-$





- consider topology in reconstruction
 - allow for flight distance for on-shell neutrinos
 - common vertex for virtual neutrinos





virtual Majorana neutrinos

 $B^- \rightarrow D^+ \mu^- \mu^-$  $B^- \rightarrow D^{*+} \mu^- \mu^-$ 

$$\mathcal{B}(B^- \rightarrow D^+ \mu^- \mu^-) < 6.9 \times 10^{-7}$$

@ 95 % CL

$$\mathcal{B}(B^- \rightarrow D^{*+} \mu^- \mu^-) < 2.4 \times 10^{-6}$$

@ 95 % CL



LHCb
FNAL

 0.41 fb^{-1} [arXiv:1201.5600](https://arxiv.org/abs/1201.5600)



on-shell Majorana neutrinos

- mis-identification rates from data with mass shape from simulation

 $B^- \rightarrow \pi^+ \mu^- \mu^-$  $B^- \rightarrow D_s^+ \mu^- \mu^-$ 

assuming B^- phase space decay:

$$\mathcal{B}(B^- \rightarrow \pi^+ \mu^- \mu^-) < 1.3 \times 10^{-8}$$

@ 95 % CL

$$\mathcal{B}(B^- \rightarrow D_s \mu^- \mu^-) < 5.8 \times 10^{-7}$$

@ 95 % CL



0.41 fb^{-1}

arXiv:1201.5600







implications on Majorana mass

mass extraction

Determine limit as function of $h^+ \mu^-$ mass

$$B^- \rightarrow \pi^+ \mu^- \mu^-$$

$$B^- \rightarrow D_s^+ \mu^- \mu^-$$



0.41 fb⁻¹

arXiv:1201.5600







new bounds for on shell Majorana neutrinos

Status of 2009

LHCb limit on $|V_{\mu 4}|$



arXiv:1201.5600




arXiv:0901.3589







$$B^- \rightarrow D^0 \pi^+ \mu^- \mu^-$$



- Four body B decay complementary to three body decay
([arXiv:1108.6009](https://arxiv.org/abs/1108.6009))
- $m_4 = m(\pi^+ \mu^-)$
 - + enhanced by W couplings
 - smaller mass range accessible ($260 \text{ MeV} < m_4 < 3.3 \text{ GeV}$)
- first performed at LHCb



$$B^- \rightarrow D^0 \pi^+ \mu^- \mu^-$$



$$\mathcal{B}(B^- \rightarrow D^0 \pi^+ \mu^- \mu^-) < 1.5 \times 10^{-6} \text{ @95 \% CL } ^a$$

Less restrictive than $B^- \rightarrow \pi^+ \mu^- \mu^-$ on $|V_{\mu 4}|^2$



^aLHCb, CERN-PH-EP-2012-006, [arXiv:1201.5600](https://arxiv.org/abs/1201.5600)



Summary on LNV in B decays

channel	limit	
$\mathcal{B}(B^- \rightarrow \pi^+ e^- e^-)$	$< 2.3 \times 10^{-8}$	@90 % CL  ^a
$\mathcal{B}(B^- \rightarrow K^+ e^- e^-)$	$< 3.0 \times 10^{-8}$	@90 % CL  ^a
$\mathcal{B}(B^- \rightarrow K^{*+} e^- e^-)$	$< 2.8 \times 10^{-6}$	@90 % CL  ^b
$\mathcal{B}(B^- \rightarrow \rho^+ e^- e^-)$	$< 2.6 \times 10^{-6}$	@90 % CL  ^b
$\mathcal{B}(B^- \rightarrow D^+ e^- e^-)$	$< 2.6 \times 10^{-6}$	@90 % CL  ^c
$\mathcal{B}(B^- \rightarrow D^+ e^- \mu^-)$	$< 1.8 \times 10^{-6}$	@90 % CL  ^c
$\mathcal{B}(B^- \rightarrow \pi^+ \mu^- \mu^-)$	$< 1.3 \times 10^{-8}$	@95 % CL  ^d
$\mathcal{B}(B^- \rightarrow K^+ \mu^- \mu^-)$	$< 5.4 \times 10^{-7}$	@95 % CL  ^e
$\mathcal{B}(B^- \rightarrow D^+ \mu^- \mu^-)$	$< 6.9 \times 10^{-7}$	@95 % CL  ^d
$\mathcal{B}(B^- \rightarrow D^{*+} \mu^- \mu^-)$	$< 2.4 \times 10^{-6}$	@95 % CL  ^d
$\mathcal{B}(B^- \rightarrow D_s^+ \mu^- \mu^-)$	$< 5.8 \times 10^{-7}$	@95 % CL  ^d
$\mathcal{B}(B^- \rightarrow D^0 \pi^- \mu^- \mu^-)$	$< 1.5 \times 10^{-6}$	@95 % CL  ^d

^aBaBar, [Phys. Rev. D 85, 071103](#) (2012)

^bCLEO, [Phys. Rev. D 65, 111102](#) (2002)

^cBelle, [Phys. Rev. D 84, 071106\(R\)](#), (2011)

^dLHCb, CERN-PH-EP-2012-006, [arXiv:1201.5600](#) (2012)

^eLHCb, [Phys. Rev. Lett. 108 101601](#) (2012)



Lepton Flavour Violation



- $D^- \rightarrow h^- \ell^+ \ell'^-$
- $B^- \rightarrow \pi^- / K^- \tau^\pm \ell^\mp$
- $\tau^- \rightarrow \mu^- \mu^- \mu^+$



Lepton Flavour Violation

Standard Model + neutrino oscillation

- established for neutrinos
- can enter charged sector in loops
- predicted rates unmeasurable small






Lepton Flavour Violation

- established for neutrinos
- can enter charged sector in loops
- predicted rates unmeasurable small
- enhancement predicted in many New Physics models, e.g.
 - multi-Higgs extensions¹
 - leptoquarks²
 - low scale seesaw models³

extended Higgs

$$D^+ \rightarrow \pi^+ \mu^- e^+$$




¹Phys. Rev. D **44**, 1461

²Z. Phys. C **61**, 613

³Phys. Rev. D **73**, 074011



LFV in charm decays




[HFAG]





LFV in bottom decays



- LFV τ sector not probed by D decays
- enough energy available in B decays
- multi Higgs models enhance $B \rightarrow h\tau\mu$ over other channels⁴
- challenging: neutrino in τ decay similar to $B \rightarrow X\ell\nu$

⁴Phys. Rev. D 44 1461



Lfv in bottom decays

channel	limit	
$\mathcal{B}(B^+ \rightarrow K^+ \tau^- \mu^+)$	$< 4.5 \times 10^{-5}$	a
$\mathcal{B}(B^+ \rightarrow K^+ \tau^+ \mu^-)$	$< 2.8 \times 10^{-5}$	a
$\mathcal{B}(B^+ \rightarrow K^+ \tau^- e^+)$	$< 4.3 \times 10^{-5}$	a
$\mathcal{B}(B^+ \rightarrow K^+ \tau^+ e^-)$	$< 1.5 \times 10^{-5}$	a
$\mathcal{B}(B^+ \rightarrow \pi^+ \tau^- \mu^+)$	$< 6.2 \times 10^{-5}$	a
$\mathcal{B}(B^+ \rightarrow \pi^+ \tau^+ \mu^-)$	$< 4.5 \times 10^{-5}$	a
$\mathcal{B}(B^+ \rightarrow \pi^+ \tau^- e^+)$	$< 7.4 \times 10^{-5}$	a
$\mathcal{B}(B^+ \rightarrow \pi^+ \tau^+ e^-)$	$< 2.0 \times 10^{-5}$	a
$\mathcal{B}(B^+ \rightarrow \pi^+ \mu^\pm e^\mp)$	$< 1.7 \times 10^{-7}$	b
$\mathcal{B}(B^+ \rightarrow K^+ \mu^+ e^-)$	$< 9.1 \times 10^{-8}$	b

details in talk of Giovanni Marchiori

LFV energy scale

$\Lambda_{\bar{b}d}$:

2.2 TeV \rightarrow 11 TeV

$\Lambda_{\bar{b}s}$:

2.6 TeV \rightarrow 15 TeV

a: BaBar, [arXiv:1204.2852](https://arxiv.org/abs/1204.2852), 429 fb $^{-1}$



b: BaBar, [Phys. Rev. Lett. 99, 051801](https://doi.org/10.1103/PhysRevLett.99.051801) (2007), 209 fb $^{-1}$





$\tau^- \rightarrow \mu^- \mu^- \mu^+$ at hadron colliders

- in Standard Model + neutrino oscillation introduced in penguin diagrams
- highly suppressed ($\mathcal{B} \ll 10^{-50}$)
- enhanced by e.g. charged Higgs
- best measurements conducted at B factories (Francesco Renga's talk)
- ! also possible at hadron colliders





$\tau \rightarrow \mu\mu\mu$ in hadron collisions

τ sources at B factories

- $e^+e^- \rightarrow \tau^+\tau^-$
- allows τ tag

τ sources at the LHC

- inclusive τ cross section $79.5 \pm 8.3 \text{ }\mu\text{b}$ at $\sqrt{s} = 7 \text{ TeV}$ in the LHCb acceptance
→ $7.9 \times 10^{10} \tau$ produced
- dominant: $D_s \rightarrow \tau\nu_\tau$ (78%)
! no τ tag possible
- standard τ signatures very similar to D decays
 $(D^+ \rightarrow \pi^+\pi^-\pi^+\pi^0$
vs. $\tau^+ \rightarrow \pi^+\pi^-\pi^+\bar{\nu}_\tau)$




Strategy

- Loose cut based selection
- Classification in 3D space:
 - invariant mass
 - decay topology (multivariate)
 - particle identification (multivariate)
- Classifier trained on simulation
- Calibration with control channel
- Normalisation with D_s
- CLs method to extract the result

LHCb
FPCP

Signal & Calibration channel






Strategy

- Loose cut based selection
- Classification in 3D space:
 - invariant mass
 - decay topology (multivariate)
 - particle identification (multivariate)
- Classifier trained on simulation
- Calibration with control channel
- Normalisation with D_s
- CLs method to extract the result

LHCb
~~FPCP~~

Signal & peaking background





Normalisation channel $D_s^+ \rightarrow \phi(\mu^+\mu^-)\pi^+$

produced τ leptons


$$\mathcal{B}(\tau \rightarrow \mu\mu\mu) = \frac{\sigma(pp \rightarrow D_s \rightarrow \tau)}{\sigma(pp \rightarrow \tau)} \frac{\mathcal{B}(D_s \rightarrow \phi(\mu\mu)\pi)}{\mathcal{B}(D_s \rightarrow \tau\nu_\tau)} \frac{\varepsilon_{norm}}{\varepsilon_{sig}} \frac{N_{\tau \rightarrow \mu\mu\mu}}{N_{D_s \rightarrow \phi(\mu\mu)\pi}}$$

ingredients

- $b\bar{b}$ and $c\bar{c}$ production measurements from LHCb
- $B \rightarrow \tau$ and $D \rightarrow \tau$ branching ratios from LEP / B factories
- $\mu\mu\pi$ final state
- systematics cancel
 - trigger
 - reconstruction
 - selection



LHCb-CONF-2012-015



45 515 $D_s \rightarrow \phi(\mu^+\mu^-)\pi$ decays





Normalisation channel $D_s^+ \rightarrow \phi(\mu^+\mu^-)\pi^+$

produced τ leptons


$$\mathcal{B}(\tau \rightarrow \mu\mu\mu) = \frac{\sigma(pp \rightarrow D_s \rightarrow \tau)}{\sigma(pp \rightarrow \tau)} \frac{\mathcal{B}(D_s \rightarrow \phi(\mu\mu)\pi)}{\mathcal{B}(D_s \rightarrow \tau\nu_\tau)} \frac{\varepsilon_{norm}}{\varepsilon_{sig}} \frac{N_{\tau \rightarrow \mu\mu\mu}}{N_{D_s \rightarrow \phi(\mu\mu)\pi}}$$

ingredients

- $b\bar{b}$ and $c\bar{c}$ production measurements from LHCb
- $B \rightarrow \tau$ and $D \rightarrow \tau$ branching ratios from LEP / B factories
- $\mu\mu\pi$ final state
→ systematics cancel
- non resonant contribution
 $< 2\%$



LHCb-CONF-2012-015



background substracted $\mu^+\mu^-$ mass





signal likelihoods


3 body decay likelihood

- vertex properties
 - vertex fit, pointing
- track quality
- isolation


particle identification

- hits in muon chambers
- energy in calorimeters
 - compatible with MIP
- RICH information

Calibration

 $D_s \rightarrow \phi\pi$ 

Calibration

 $J/\psi \rightarrow \mu^+\mu^-$ 



invariant mass


two fold role

- background estimation in sidebands
- different signal likelihood inside signal region

LHCb
TFCP

1 fb^{-1}

LHCb-CONF-
2012-015



- mass resolution and mass scale calibrated on data






signal likelihoods

combined signal distribution

- events distributed over 25 likelihood bins
- background estimate from mass sidebands


LHCb
FPCP 1 fb^{-1}
LHCb-CONF-
2012-015





- 11 % signal efficiency
- 21 % signal efficiency
- for illustration: high likelihood range shown




observed events



11 % of the signal
0.03 % of the background



21 % of the signal
0.14 % of the background



red dashed combinatorial background
green $D_s^+ \rightarrow \eta(\mu^-\mu^+\gamma)\mu^+\nu_\mu$
blue combined background



1 fb^{-1}

LHCb-CONF-2012-015





extracted limit


result (preliminary)

	observed	expected
$\mathcal{B}(\tau^- \rightarrow \mu^- \mu^- \mu^+)$	$< 6.3 \times 10^{-8}$	8.2×10^{-8} @ 90 % CL
	$< 7.8 \times 10^{-8}$	9.9×10^{-8} @ 95 % CL

LHCb
FCCP

1 fb⁻¹

LHCb-CONF-
2012-015



Belle 2.1×10^{-8} @ 90 % CL

Phys. Lett. B **687**, 139
(2010)

BaBar 3.3×10^{-8} @ 90 % CL

Phys. Rev. D **81**,
111101(R) (2010)





Summary

- neither lepton number violation nor lepton flavour violation observed (yet)
 - hadron decays provide good probes for Lepton Number & Flavour Violation
 - excellent sensitivity achieved at B factories
 - LNV and LFV excluded to 10^{-5} for D decays
 - LHCb extends knowledge on forbidden B decays
 - LNV excluded to $10^{-6\cdots -8}$
 - LHCb approaching B factories for $\tau \rightarrow \mu\mu\mu$
- ⇒ first time done at hadron colliders



BACKUP





Sensitivity to long lived Majorana neutrinos

- analysis allows for neutrino flight distance
- probes only Majorana neutrinos decaying inside the detector
- sensitivity to lifetime provided

a) $B^- \rightarrow \pi^+ \mu^- \mu^-$


$250 \text{ MeV} < m_4 < 5 \text{ GeV}$

b) $B^- \rightarrow D_s^+ \mu^- \mu^-$

$2.1 \text{ GeV} < m_4 < 5.15 \text{ GeV}$

c) $B^- \rightarrow D^0 \pi^+ \mu^- \mu^-$

$260 \text{ MeV} < m_4 < 3.3 \text{ GeV}$



- sensitive to lifetimes $< 10^{-10} \text{ s}$
- $B^- \rightarrow D^0 \pi^+ \mu^- \mu^-$ better sensitive to “large” lifetimes





model dependence

$$\tau^- \rightarrow \mu^-\mu^-\mu^+$$

possible problem

if the selection efficiency is different for various models, the limit could be higher or lower than the one we quote.

crosschecks

We tested the

- trigger efficiency
- preselection
- multivariate classifier

as a function of the invariant dimuon mass(es).

⇒ these are constant within 15 %





Prospects

When will LHCb catch up with the B factories on $\tau^- \rightarrow \mu^-\mu^-\mu^+$?

short answer

not studied so far

long answer


Judge yourself:

- used: 1 fb^{-1} @ 7 TeV
- 2012: 1.5 fb^{-1} @ 8 TeV (charm cross section $\sim 15\%$ higher)
- 2013-2014: long shutdown 1
- 2014-2017: running at higher energies



HFAG limits for bottom decays

Lepton Number Violating Charmless B Decays



[HFAG]

