

**Search for the suppressed  
 $\Lambda_c^+ \rightarrow p\mu^+\mu^-$  decay and  
observation of the  $\Lambda_c^+ \rightarrow p\omega$   
decay**



*Marcin Chrzaszcz*  
*mchrzaszcz@cern.ch*



With M. Jezabek, T. Lesiak, B. Nowak, M. Witek (IFJ PAN)

Tuesday meeting, CERN  
September 26, 2017

# Yellow pages

- ⇒ Reviewers: Tom Blake(chair), Harry Cliff, Simon Eydelman(EB)
- ⇒ Twiki:  
<https://twiki.cern.ch/twiki/bin/viewauth/LHCbPhysics/Lc2PMuMu>
- ⇒ Review start: 01.04.2017
- ⇒ Three interactions with the review committee.
- ⇒ Unbinding: 18.07.2017
- ⇒ Minor changes to the analysis during the review.

We would like to take this occasion and than Tom, Harry and Simon for fruitful, constructive and smooth review!

# Motivation

⇒ SM predictions:




$$\mathcal{O}(10^{-8})$$

⇒ Long distance effects:

$$\mathcal{O}(10^{-6})$$

⇒ Previous measurement done by Babar:

$$\text{Br}(\Lambda_c^+ \rightarrow p\mu^+\mu^-) < 4.4 \cdot 10^{-5} \text{ at } 90\% \text{ CL}$$




Should be able to improve by a factor of 100!

# Analysis strategy

- ⇒ Normalization to  $\Lambda_c^+ \rightarrow p\phi(\mu\mu)$ .
- ⇒ Typical steps rare decays:
  - Loose stripping selection.
  - BDT1 used for first preselection.
  - BDT2 used to further suppress the background.
  - PID used to fight the peaking background.
- ⇒ Search performed in several dimuon mass windows.
- ⇒ Selection optimized on  $CL_s$ .
- ⇒ Unblinding and calculate the UL of BR using  $CL_s$ .

# Analysis strategy

- ⇒ Normalization to  $\Lambda_c^+ \rightarrow p\phi(\mu\mu)$ .
- ⇒ Typical steps rare decays:
  - Loose stripping selection.
  - BDT1 used for first preselection.
  - BDT2 used to further suppress the background.
  - PID used to fight the peaking background.
- ⇒ Search performed in several dimuon mass windows.
- ⇒ Selection optimized on  $CL_s$ .
- ⇒ Unblinding and calculate the UL of BR using  $CL_s$ .



## Normalization channel

Use the  $\Lambda_c^+ \rightarrow p\phi(\mu\mu)$ .

⇒ Same final state, same selection, a lot of systematics cancel.

⇒ The Branching fraction of  $\Lambda_c^+ \rightarrow p\phi$  is known with 22 %.

Use the  $\Lambda_c^+ \rightarrow pK\pi$ .

⇒ Precisely known branching fraction (precision: 6.4 %).

⇒ A lot of additional systematics due to different final states, different selections



We choose the  $\Lambda_c^+ \rightarrow p\phi(\mu\mu)$  option

⇒ In the most optimistic scenario where you assume the 22 % systematic to go down to 6.4 % the UL.

⇒ In this case the UL gets worse 7.8 %.

# Normalization channel

Use the  $\Lambda_c^+ \rightarrow p\phi(\mu\mu)$ .

⇒ Same final state, same selection, a lot of systematics cancel.

⇒ The Branching fraction of  $\Lambda_c^+ \rightarrow p\phi$  is known with 22 %.

Use the  $\Lambda_c^+ \rightarrow pK\pi$ .

⇒ Precisely known branching fraction (precision: 6.4 %).

⇒ A lot of additional systematics due to different final states, different selections



We choose the  $\Lambda_c^+ \rightarrow p\phi(\mu\mu)$  option

⇒ In the most optimistic scenario where you assume the 22 % systematic to go down to 6.4 % the UL.

⇒ In this case the UL gets worse 7.8 %.

# Data sets and Stripping

⇒ 2011+2012 (aka Run1) Stripping 20.

Condition	$\Lambda_c^+ \rightarrow p\mu^+\mu^-$
$\mu^\pm$ and $p$ $p_T$ Track $\chi^2/\text{ndf}$ IP $\chi^2/\text{ndf}$ PID $\mu^\pm$ PID $p$	$> 300\text{MeV}/c$ $< 3$ $> 9$ PIDmu $> -5$ and (PIDmu - PIDK) $> 0$ PIDp $> 10$
$\Lambda_c^+$ $\Delta m$ Vertex $\chi^2$ IP $\chi^2$ $c\tau$ Lifetime fit $\chi^2$	$< 150\text{MeV}/c^2$ $< 15$ $< 225$ $> 100\mu\text{m}$ $< 225$



# Preselection

⇒ Additional cuts:

Common cuts
$m_{\mu\mu} < 1400 \text{ MeV}/c^2$
proton $ProbNNp > 0.1$
$\mu^+, \mu^-$ $ProbNNmu > 0.1$
$10 \text{ GeV}/c < p_{proton} < 100 \text{ GeV}/c$

⇒ We define couple of dimuon mass regions:

$m(\mu\mu)$ region	$\text{MeV}/c^2$
$\phi$ region	[985, 1055]
$\omega$ region	[759, 805]
<i>non resonant</i>	[210, 747] $\cup$ [817, 980] $\cup$ [1060, 1400]

# Trigger

⇒ We require the following triggers (all are TOS):

- Lo
  - LoMuonDecision
- HLT1
  - Hlt1TrackMuonDecision
  - Hlt1DiMuonLowMassDecision
  - Hlt1TrackAllLoDecision
- HLT2
  - Hlt2DiMuonDetachedDecision
  - Hlt2CharmSemilep3bodyD2KMuMuDecision
  - Hlt2CharmSemilepD2HMuMuDecision

⇒ The TIS increase the signal yield by  $< 10\%$  and were asked to be removed at the WG review stage.




# BDT1 training

⇒ The normalization channel is also a rather “rare decay”:

$$\text{Br}(\Lambda_c^+ \rightarrow p\phi) \cdot \text{Br}(\phi \rightarrow \mu\mu) = 3.1 \cdot 10^{-7}$$

⇒ After the previous preselection a simple BDT is trained using variables that are well simulated in the MC. k-folding used ( $k = 10$ )



⇒ The BDT1 (not surprisingly) likes the prompt  $\Lambda_c$  rather the secondary ones.



# BDT1 selection

- ⇒ The selection based on BDT1 is not optimised.
- ⇒ A loose cut:

$$\text{BDT1} > -0.1$$





- ⇒ The normalization channel peak is observed.

# BDT2 selection

⇒ Variables used:




- flight distance - the one between the production and decay points.
- $\chi^2$  of flight distance,
- transformed decay time -  
 $T = \exp(-1000 \cdot \tau/\text{ns})$ ,
- IP - impact parameter with respect to primary vertex,
- $\chi^2$  of IP of  $\Lambda_c^+$
- $\log(\chi_{DTF}^2)$ ,
- $p_T$  - transverse momentum of  $\Lambda_c^+$ ,
- minimum of  $\chi^2$  of  $p, \mu^+, \mu^-$  w.r.t. primary vertex,
- transverse momenta



# BDT2

⇒ After correcting the DATA/MC differences the BDT distribution shows a good DATA/MC agreement.




⇒ No mass correlation observed.



# PID

⇒ MC re sampling is choose to correct the PID distributions:  
For MC samples the ProbNNp and ProbNNmu are drawn from the PIDCalib distributions.






- ⇒ The PIDCalib doesn't cover to low  $p_T$  muons (10%).
- ⇒ Decided to use for them the  $D_s \rightarrow \phi(\mu\mu)$  sample.



# Selection optimization

- ⇒ The final selection of the analysis is optimized!
- ⇒  $CL_s$  method used.
- ⇒ KDE used to sample toy experiments.

Variable	Condition
BDT	$> 0.0$
$ProbNNp(p)$	$> 0.68$
minimum $ProbNNmu(\mu^\pm)$	$> 0.38$











# Peaking backgrounds

⇒ The tight PID cuts essentially kill the peaking bkg!

⇒ The only bkg left is the  $\Lambda_c^+ \rightarrow p\pi\pi$ .



⇒ Estimated contamination:  
 $1.96 \pm 1.13$  ⇒ assigned as systematic



# Normalization

⇒ The gold equation:


$$\frac{\mathcal{B}(\Lambda_c^+ \rightarrow p\mu^+\mu^-)}{\mathcal{B}(\Lambda_c^+ \rightarrow p\phi(\mu\mu))} = \frac{\epsilon_{\text{norm}}^{\text{TOT}}}{\epsilon_{\text{sig}}^{\text{TOT}}} \times \frac{N_{\text{sig}}}{N_{\text{norm}}},$$

⇒ We take advantage of the cancellation that:

$$\frac{\epsilon_{\text{norm}}^{\text{TOT}}}{\epsilon_{\text{sig}}^{\text{TOT}}} = \frac{\epsilon_{\text{norm}}^{\text{STRIP}}}{\epsilon_{\text{sig}}^{\text{STRIP}}} \times \frac{\epsilon_{\text{norm}}^{\text{COMM}}}{\epsilon_{\text{sig}}^{\text{COMM}}} \times \frac{\epsilon_{\text{norm}}^{\text{SPEC}}}{\epsilon_{\text{sig}}^{\text{SPEC}}}, \quad \frac{\epsilon_{\text{norm}}^i}{\epsilon_{\text{sig}}^i} \simeq 1$$


⇒ In addition we have added 6 mass bins to increase the sensitivity.

⇒ Signal is modelled by a double Gaussian.





# Expected background

⇒ Background modelled with a linear function.



bin	no events
bin1	$8.56136 \pm 0.540302$
bin2	$8.60318 \pm 0.536917$
bin3	$8.64582 \pm 0.536561$
bin4	$8.6887 \pm 0.539208$
bin5	$8.7304 \pm 0.544752$
bin6	$8.77226 \pm 0.553162$






⇒ Expected upper limits:  
 $\mathcal{B}(\Lambda_c^+ \rightarrow p\mu^+\mu^-) <$   
 $5.91 \times 10^{-8}$  at 90 % CL

# Observed Upper limits

⇒ After the green light from RC we have unblinded we did not observed a significant excess of events.

⇒ We have set an UL:

$$\mathcal{B}(\Lambda_c^+ \rightarrow p\mu^+\mu^-) < 7.68 \times 10^{-8} \text{ at } 90\% \text{ CL}$$



# By product :)



⇒ We also looked at the  $\omega$  dimuon region.

We observed an excess

Using Wilks theorem we have calculated the significance to be  $5.0 \sigma$ !

⇒ This is the first observation of this decay!!!

$$\mathcal{B}(\Lambda_c^+ \rightarrow p\omega) = (7.6 \pm 2.6 \text{ (stat)} \pm 0.9 \text{ (syst1)} \pm 3.1 \text{ (syst2)}) \times 10^{-4}$$




# Conclusion

- Improved the UL for  $\mathcal{B}(\Lambda_c^+ \rightarrow p\mu^+\mu^-)$  by two orders of magnitude!

# Conclusion

- Improved the UL for  $\mathcal{B}(\Lambda_c^+ \rightarrow p\mu^+\mu^-)$  by two orders of magnitude!



- First time observed the decay  $\Lambda_c^+ \rightarrow p\omega$ !!
- Paper is being prepared, aiming PRL
- We would like to ask the collaboration for approving this analysis.

