

# Low Mass Drell-Yan Status Report



Marcin Chrząszcz  
Katharina Müeller  
Nicola Chiapolini





University of  
Zurich <sup>UZH</sup>

Analysis and software week, CERN  
January 27, 2015

# Introduction to Drell-Yan

- Drell-Yan are process of two quark annihilations in which neutral coupling to two leptons.
- The cross section of this process depends on two components:
  - Hard scattering process  $\Rightarrow$  NNLO pQCD.
  - Parton Distribution Function (PDF).
- Measurement of the cross section have a high sensitivity to the PDF
- Due to unique coverage  $2 < y < 5$  LHCb probes the  $Q^2 - x$  region not covered by other experiments.



# Selection

- Analysis based on 2011 and 2012 data set.
- Plan to measure them separately as well as the ratio (cancellation of systematics).
- Trigger:
  - L0\_L0DiMuonDecision,
  - Hlt1DiMuonHighMassDecision,
  - Hlt2DiMuonDY(3,4)Decision
- Stripping:
  - StrippingDY2MuMuLine(3,4)
- Selection:
  - $2 < \eta^\mu < 4.5$ ,
  - $p^\mu > 10 \text{ GeV}$ ,
  - $p_T^\mu > 3 \text{ GeV}$ ,
  - $\chi_{vtx}^{2,\mu\mu} < 5$ ,
  - $10 < m(\mu\mu) < 120 \text{ GeV}$ .

# The Goal

⇒ Since there is no normalization channel, we will use the integrated luminosity for cross section calculations

⇒ The measurement will be performed in the bins of dimuon mass and pseudo-rapidity:

	10.5 – 11.0	11.0 – 11.5	11.5 – 12.0
	12.0 – 13.0	13.0 – 14.0	14.0 – 15.0
$M_{\mu\mu}$ [GeV/ $c^2$ ]	15.0 – 17.5	17.5 – 20.0	20.0 – 25.0
	25.0 – 30.0	30.0 – 40.0	40.0 – 60.0
	60.0 – 70.0	70.0 – 80.0	80.0 – 90.0
	90.0 – 100.0	100.0 – 110.0	110.0 – 120.0
$y$	2.0 – 4.5		

$M_{\mu\mu}$ [GeV/ $c^2$ ]	10.5 – 12.0	12.0 – 15.0	
	15.0 – 20.0	20.0 – 60.0	
$y$	2.0 – 2.25	2.25 – 2.5	2.5 – 2.75
	2.75 – 3.0	3.0 – 3.25	3.25 – 3.5
	3.5 – 4.5		



# Isolation

- Drell-Yan unfortunately do not peak in mass  $\rightarrow$  need another variable to control the purity.
- Instead we define an isolation variable:

$$\mu_{\text{iso}} = \log(p_T^{\text{cone}}(\mu, 0.5) - p_T^{\text{cone}}(\mu, 0.1))$$



- For two muons we take the maximum of the two isolations:

$$\mu\mu_{\text{iso}} = \max(\mu_{\text{iso}}^+, \mu_{\text{iso}}^-)$$





# Isolation mass dependence

- Unfortunately the  $\mu\mu_{iso}$  is showing some mass dependence:



(a) data





(b) simulation

# Signal template

- We do not want to use MC for determination of the signal  $\mu\mu_{iso}$  template.
- We adopted a data driven procedure:
  - The template is taken from data and scaled to account for  $\mu\mu_{iso}$  mass dependence.
- Possibility 1:
  - Take the  $Splot Z \rightarrow \mu\mu$  from data and multiply it by the scale factor determined from minimalising the  $\chi^2$  between MC  $Z$  and DY in particular region.

A RooPlot of "Z0\_MM"



# Signal template



- Possibility 2:

- Use a second decay from data:  $\Upsilon \rightarrow \mu\mu$ .
- The template for a given mass range ( $M_{\min}, M_{\max}$ ) is choose as:

$$\text{Temp}(M) = \text{Temp}^{\Upsilon} \frac{(M_Z - M_{\Upsilon} - (M - M_{\Upsilon}))}{M_Z - M_{\Upsilon}} \\ + \text{Temp}^Z \frac{M - M_{\Upsilon}}{M_Z - M_{\Upsilon}}$$

- Then the new obtained template is scaled in the same way as the previous one.


A RooPlot of "Z0\_MM"





# Signal template - Summary

- We are investigating the impact on the analysis for the different approaches
- For now it looks like the results do not change with using different signal templates.
- Because templates are data driven we need to ensure a large statistics in each of the  $m_{\mu\mu}$ ,  $y$  bins, because of this the last  $y$  bin is larger then the rest.



# Backgrounds


- There are two sources of backgrounds:
  - Heavy flavour decays.
  - Mis-ID.
- For fitting the  $\mu\mu_{iso}$  we need to know both the signal and background distribution.
- Background templates can be determined from data
  - Heavy flavour decays:
    - ↔ Requiring the  $\chi_{vtx}^{2,\mu\mu} > 16$
    - ↔ For cross-check IP  $> 5$  mm
  - Miss-ID:
    - ↔ Require that both muons have the same sign.
    - ↔ For cross-check take the minimum bias stripping line.

# Over all fits


- Using the above 3 mentioned templates the fits converge without any problems.
- The higher one goes in mass the cleaner the signal is.

Mass bin	Purity
[40, 60] GeV	$0.879 \pm 0.019$
[30, 40] GeV	$0.754 \pm 0.015$
[25, 30] GeV	$0.657 \pm 0.011$
[20, 25] GeV	$0.507 \pm 0.008$
[17.5, 20] GeV	$0.402 \pm 0.007$
[15, 17.5] GeV	$0.316 \pm 0.006$

Data\_12000\_15000\_y\_bin\_2\_2.25



Data\_15000\_20000\_y\_bin\_3.5\_4.5



# Cross section calculations

- To calculate the cross section the luminosity will be used:

$$\sigma = \frac{\rho f^{\text{MIG}}}{\mathcal{L} \epsilon^{\text{SEL}}} \sum \frac{1}{\epsilon^{\text{TRIG}} \epsilon^{\text{MUID}} \epsilon^{\text{GEC}} \epsilon^{\text{TRACK}}},$$

where



- $\rho$  signal fraction from the fit.
- $f^{\text{MIG}}$  correction to bin-bin migration.
- $\mathcal{L}$  integrated luminosity.
- $\epsilon^{\text{SEL}}$  efficiency on the vertex requirement.
- $\epsilon^{\text{MUID}}$  muon identification efficiency.
- $\epsilon^{\text{GEC}}$  global event cut efficiency.
- $\epsilon^{\text{TRACK}}$  tracking efficiency.

# Luminosity

- Thanks to our colleagues the error on the luminosity in LHCb is 1.16(1.71)% for 2012(2011) data.
- For the 8 TeV data we removed: 111802-111890 , 126124-126160, 129530-129539 runs.
- Lost 14.68 pb<sup>-1</sup> of data in total.
- For the 7 TeV data we removed: 101401, 101403-101415 runs.
- Lost 8.23 pb<sup>-1</sup>.

# Trigger efficiency


- We take the trigger efficiency from MC. We are using the dimuon trigger that were always well simulated.
- We performed a cross check using tag and probe method that ensures the luminosity is correctly simulated.




- An systematic uncertainty of 0.01 is assigned.

# Muon Identification



- Only muon ID requirement in this analysis is the `isMuon`.
- The efficiency is taken from MC.
- Has been cross-checked that it agrees in LHCb-INT-2014-030



(a)  $\mu^-$




(b)  $\mu^+$



- The systematics is 0.005 (needs to be checked for the low  $p_T$ ).

# Global even cut efficiency

- There is a SPD cut for the dimuon trigger:  $\text{SPD} < 900$ .
- A data driven method is used to estimate the cut.



- No dependence is observed of the  $M_{\mu\mu}$  and the  $y$  in data.
- Similar to the  $W$  and  $Z$  analysis.



# Conclusions

- Analysis is well advanced!
- The analysis note is being written as we speak:  
svn+ssh:  
`//svn.cern.ch/repos/lhcbdocs/Users/mchrzasz/DY_ANANote`
- +30 pages!
- To do list:
  - Calculate the theory predictions for 8 TeV data.
  - Missing systematics: bin-bin migration, templates determination.
  - Hopefully the ANA note in WG review soon!

# Backup