

# RECO 12 vs 14, vol 1

Tom Blake<sup>1</sup>, Marcin Chrzęszcz<sup>2,3</sup>

<sup>1</sup> University of Warwick,

<sup>2</sup> University of Zurich,

<sup>3</sup> Institute of Nuclear Physics, Krakow



**University of  
Zurich**<sup>UZH</sup>



August 18, 2014





As you should now from email exchange:

- $B^0 \rightarrow K^* \mu\mu$  PHPS MC was produced.
- We also have SM MC.
- $J/\psi K^*$  on way(later today we should have it).
- All are SIM08
- For now 4  $TeV$  data, 3.5  $TeV$  in queue.
- The same events(common, Gauss, Boole, Moore) are processed by 2 different Brunel versions(one for RECO12, other for RECO14).
- Since I don't have full selection, I am studying the events that pass our stripping 20.
- Events are truth matched between two ntuples using EVENTNUMBER.




- RECO 12 and STRIPPING 20 selected 5054
- Among those 4667 are also selected by RECO 14 and our stripping.
- The over lap is about  $92.3 \pm 0.4\%$ .
- A bit higher compared to what we saw in data....

# Results/Plots $B^0$




# Results/Plots B<sup>0</sup>


B<sub>0</sub>\_IPCH12




B<sub>0</sub>\_MM



B<sub>0</sub>\_PT




B<sub>0</sub>\_PT




# Results/Plots $B^0$




B0\_tau






B0\_VRTCHI2






# Results/Plots K











# Results/Plots $\mu^+$


Mup\_P



Mup\_PT



Mup\_TRK\_CHI2



# Conclusions

- 1 The spread of the variables looks smaller than in data.
- 2 This needs further investigations:  $J/\psi K^*$ , full selection, 3.5 TeV, (more suggestions?)
- 3 What else should I compare? PID? other? Let me know.
- 4 The thing that comes to my mind when I look at those plots is:

"The truth is rarely pure and never simple."